



Fairfield University
DigitalCommons@Fairfield

Mathematics Faculty Publications

Mathematics Department

2005

Markov extensions and conditionally invariant measures for certain logistic maps with small holes

Mark Demers

Fairfield University, mdemers@fairfield.edu

Follow this and additional works at: <https://digitalcommons.fairfield.edu/mathandcomputerscience-facultypubs>

Copyright 2005 Cambridge University Press

Peer Reviewed

Repository Citation

Demers, Mark, "Markov extensions and conditionally invariant measures for certain logistic maps with small holes" (2005). *Mathematics Faculty Publications*. 36.

<https://digitalcommons.fairfield.edu/mathandcomputerscience-facultypubs/36>

Published Citation

Mark Demers, "Markov extensions and conditionally invariant measures for certain logistic maps with small holes," *Ergodic Theory and Dynamical Systems* 25:4 (2005), 1139-1171.

This Article is brought to you for free and open access by the Mathematics Department at DigitalCommons@Fairfield. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of DigitalCommons@Fairfield. For more information, please contact digitalcommons@fairfield.edu.

Markov extensions and conditionally invariant measures for certain logistic maps with small holes

MARK F. DEMERS

Department of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA
(e-mail: demers@math.gatech.edu)

(Received 6 June 2004 and accepted in revised form 19 October 2004)

Abstract. We study the family of quadratic maps $f_a(x) = 1 - ax^2$ on the interval $[-1, 1]$ with $0 \leq a \leq 2$. When small holes are introduced into the system, we prove the existence of an absolutely continuous conditionally invariant measure using the method of Markov extensions. The measure has a density which is bounded away from zero and is analogous to the density for the corresponding closed system. These results establish the exponential escape rate of Lebesgue measure from the system, despite the contraction in a neighborhood of the critical point of the map. We also prove convergence of the conditionally invariant measure to the SRB measure for f_a as the size of the hole goes to zero.

1. Introduction

Consider a particle on a billiard table with convex boundaries so that the dynamics of the particle are hyperbolic, i.e. the trajectories are unstable with respect to initial conditions. Suppose a small hole is made in the table. What are the statistical properties of the trajectories in this system? If p_n is the probability that a trajectory remains on the table until time n , what is the decay rate of p_n ? More generally, we can place a particle randomly on the table according to an initial distribution μ_0 . If μ_n represents its normalized distribution at time n (assuming the particle has not escaped by time n), does μ_n converge to some μ independent of μ_0 ? Such a measure μ is a *conditionally invariant measure* for the open billiard system.

Considering the billiard table with a small hole as a perturbation of the billiard table with no holes, we can pose a related question in terms of the stability of the closed system: does the conditionally invariant measure of the open system converge to the invariant measure of the closed system as the size of the hole tends to zero?

The billiard table with a hole as a model for an open chaotic dynamical system was proposed by Pianigiani and Yorke [PY]. Although these questions remain open, dynamical systems with holes have been studied in some detail. Mathematical results

so far have focused on open systems which are uniformly hyperbolic. Pianigiani and Yorke [PY] and later Collet *et al* [CMS1, CMS2] studied expanding maps which admit a finite Markov partition after the introduction of holes. These results were generalized to smooth Smale horseshoes [C1, C2] and a class of scattering billiards with a non-eclipsing condition [LM, R]. Recently, Chernov and Markarian [CM2, CM1] studied Anosov diffeomorphisms with holes which were elements of a finite Markov partition. In [CMT1] and [CMT2], the Markov restriction on the holes was relaxed, but the results still used strongly the Markov partitions associated with Anosov diffeomorphisms.

In low-dimensional settings, efforts to drop the Markov requirements on both the map and the holes have had some success for expanding maps of the interval. A spectral analysis of the transfer operator was performed in [BK] and the stability of the spectrum was established in [KL] for perturbations of expanding maps including small holes. More constructive techniques using bounded variation and contraction mapping arguments have been used in [BCh] and [LiM] to prove the existence and properties of conditionally invariant measures. Markov extensions were used in [D] to drop some of the earlier technical requirements and limit only the size of the holes.

This brief survey highlights the classes of systems with holes which have been studied to date: expanding maps in one dimension; and in higher dimensions, systems which admit finite Markov partitions. These systems are all uniformly hyperbolic.

In this paper, we seek to understand the escape dynamics of a class of open systems which are not uniformly hyperbolic, but which do exhibit exponential recurrence times. To do this, we construct Markov extensions for certain parameter values of the logistic family after the introduction of holes. We then use the results obtained in [D] for abstract tower maps with holes to determine the existence and properties of a conditionally invariant measure.

In this use of Markov extensions, we follow the approach of Young in [Y2], in which Markov extensions were used to study a variety of closed systems including Axiom A diffeomorphisms, piecewise hyperbolic maps, Hénon maps, logistic maps, and a class of scattering billiards (see also [Y3, BY]). Chernov [Ch1, Ch2, Ch3] has also used this technique to study the statistical properties of other chaotic systems. By extending the use of Markov extensions to open systems, we hope to be able to study more general classes of systems with holes and in particular those which satisfy neither uniform hyperbolicity nor Markov requirements.

The outline of the paper is as follows. In §2 we introduce the class of logistic maps which we shall study in this paper and state our main results. Section 3 reviews the setting and main results for tower maps in [D] which we will use. In §4 we construct Markov extensions for our class of logistic maps with holes and in §5, we use the results of §3 to determine the existence and properties of a conditionally invariant measure.

1.1. Conditionally invariant measures. The problem of the billiard table with a hole can be reformulated for maps of the interval as follows.

Let \hat{T} be a map of an interval \hat{I} to itself. We take the hole H to be a finite union of open intervals and keep track of the iterates of a point until it reaches the hole. Once a point enters H , it is not allowed to return.

Let $I = \hat{I} \setminus H$ and let $T = \hat{T}|(I \cap \hat{T}^{-1}I)$. A probability measure μ on \hat{I} is said to be conditionally invariant if

$$\frac{\mu(T^{-1}A)}{\mu(T^{-1}I)} = \mu(A)$$

for every Borel subset A of \hat{I} . The measure μ is called an *absolutely continuous conditionally invariant measure* (abbreviated a.c.c.i.m.) if it is absolutely continuous with respect to Lebesgue measure.

The quantity $\lambda = \mu(T^{-1}I)$ is called the eigenvalue of the measure and $-\log \lambda$ represents the exponential rate at which mass escapes from the system. From the point of view of physical observables, we are interested in conditionally invariant measures whose escape rate indicates the rate at which (normalized) Lebesgue measure escapes from the system. For this reason, in this paper we will restrict our attention to the existence and properties of absolutely continuous conditionally invariant measures.

2. Setting and statement of results

We begin by defining the class of logistic maps that we shall study in this paper.

2.1. A class of logistic maps. One class of logistic maps which has been studied in some detail are those satisfying the Misiurewicz condition: namely, that there are no attracting or semi-attracting periodic orbits. In this paper, we study parameter values of a for which f_a satisfies a slightly more generalized set of conditions. This approach follows that of Wang and Young in [WY2]. We define the class of maps \mathcal{M} as follows.

Definition 2.1. The logistic map $f = f_a$ is in \mathcal{M} if there exists a $\delta_0 > 0$ such that the following conditions hold.

- (a) The critical orbit is bounded away from 0: $\text{dist}(f^n(0), 0) > 2\delta_0$ for all $n > 0$.
- (b) Dynamics outside of $(-\delta_0, \delta_0)$: there exist $\lambda_0 > 0$, $M_0 \in \mathbb{Z}^+$ and $0 < c_0 \leq 1$ such that:
 - (i) for all $n \geq M_0$, if $x, f(x), \dots, f^{n-1}(x) \notin (-\delta_0, \delta_0)$, then $|(f^n)'(x)| \geq e^{\lambda_0 n}$;
 - (ii) for any n , if $x, f(x), \dots, f^{n-1}(x) \notin (-\delta_0, \delta_0)$ and $f^n(x) \in (-\delta_0, \delta_0)$, then $|(f^n)'(x)| \geq c_0 e^{\lambda_0 n}$.
- (c) Recovery time for $x \in (-\delta_0, \delta_0)$: for all $x \in (-\delta_0, \delta_0)$, there exists $s_0(x) \sim \log 1/|x|$ such that $f^j(x) \notin (-\delta_0, \delta_0)$ for all $j < s_0$ and $|(f^{s_0})'(x)| \geq c_0^{-1} e^{\frac{1}{3}\lambda_0 s_0}$.

Lemma 2.5 of [WY1] implies that maps satisfying the Misiurewicz condition belong to \mathcal{M} . In viewing this class of maps, we divide the phase space into two parts: $(-\delta_0, \delta_0)$ and its complement. Part (b) of the definition says that f is essentially expanding outside of $(-\delta_0, \delta_0)$ while part (c) ensures that when orbits come close to the critical point, they subsequently spend enough time away from $(-\delta_0, \delta_0)$ for their derivatives to recover some exponential growth.

Although our method of proof will work for any map satisfying the above definition, for definiteness, we take a near 2 in the proofs contained in §4. In this parameter range, we think of λ_0 as $\log 1.9$.

2.2. Introduction of the hole. A hole H in $[-1, 1]$ is a finite union of open intervals H_j , $j = 1, \dots, L$. We wish to study the dynamics of $f_a \in \mathcal{M}$ on $[-1, 1] \setminus H$ and in particular to establish an exponential rate of escape from $[-1, 1]$. To this end, we define $\hat{I} = [-1, 1]$, $I = \hat{I} \setminus H$. We fix a , let $\hat{T} = f_a$ and set $I^n = \bigcap_{i=0}^n \hat{T}^{-i} I$. Let $T = \hat{T}|_{I^1}$.

Our first assumption on the hole involves its location in $[-1, 1]$.

Condition A1. The critical orbit is bounded away from H .

We define r to be the smaller of this distance and δ_0 .

Our second condition on H is that the positions of its components are generic with respect to one another.

Condition A2. For a fixed $m_0 \in \mathbb{Z}^+$, there exists $\varepsilon_0 > 0$ such that for any interval $\omega \subset I$, if $|\hat{T}^i \omega| < \varepsilon_0$ for $0 \leq i \leq m_0$, then there is at most one i and one j such that $\hat{T}^i \omega \cap H_j \neq \emptyset$.

We are free to choose m_0 and generally m_0 will depend on λ_0 . For a near 2, we mentioned earlier that $\lambda_0 = \log 1.9$ and in this case $m_0 = 10$ is large enough.

Practically, Condition A2 may be difficult to check. However, if we let $N_{\varepsilon_0}(A)$ be the deleted ε_0 -neighborhood of a set A , then Condition A2 is implied by the simpler condition $N_{\varepsilon_0}(\hat{T}^i H) \cap N_{\varepsilon_0}(\hat{T}^j H) = \emptyset$ for every pair $i \neq j$, $0 \leq i, j \leq m_0$.

The third assumption is on the size of the hole. We use m interchangeably as Lebesgue measure on both the tower and on the interval $[-1, 1]$. Let θ be the exponential rate of return given by Proposition 4.2, ε be the length scale of the reference intervals $\Lambda^{(i)}$ and D be the constant in Lemma 4.4. Our assumption on the measure of the hole is the following.

Condition A3.

$$m(H) < \frac{(1 - \sqrt{\theta})^3}{3} \cdot \frac{\varepsilon^2}{D\theta}.$$

As we will see in §4, $\varepsilon \sim \delta^2$ and $D \sim 1/\delta$ so the restriction on the size of the hole is $\sim \delta^5$ where $\delta < \delta_0$ defines a neighborhood of the critical point which we shall use to keep track of intervals that pass near the critical point and subsequently need time for their derivatives to recover. Condition A3 in turn implies the following upper bound

$$m(H) \leq \frac{\varepsilon}{2} \frac{e^{\frac{1}{3}\lambda_0 m_0} - 2}{e^{\frac{1}{3}\lambda_0(m_0-1)}} c'_0 \delta \quad (1)$$

which we shall use to ensure that a part of every piece of length ε is returned to the base of the tower. Condition A3 itself is used to ensure that the tower we construct satisfies Condition H2 of §3 on the measure of the holes in the tower. Equation (1) is implied by Condition A3 for small δ since it only requires that $m(H) \sim \delta^3$.

These three conditions will allow us to construct a Markov extension for T with an exponential rate of return. In order to prove that the conditionally invariant measure obtained in §5.3 is bounded away from zero, we need a transitivity condition. We shall use the following fact about maps in the class \mathcal{M} :

$$\exists n_0 = n_0(\varepsilon_0) \text{ such that for every interval } J \subseteq \hat{I} \text{ with } |J| \geq \varepsilon_0/2, \hat{T}^{n_0} J \supseteq [1 - a, 1].$$

For each component of the hole H_j , $\hat{T}^{-1}(\hat{T}H_j)$ is comprised of two intervals: H_j and its symmetric counterpart which we shall call G_j . The transitivity condition we need on the holes is stated in terms of the integer n_0 and the collection of intervals H_j and G_j .

Condition A4.

- (a) $\hat{T}^i H_j \cap G_k = \emptyset$ for all $j, k \in [1, \dots, L]$, $0 \leq i \leq n_0$.
- (b) $\hat{T}^i H_j \cap H_k = \emptyset$ for all $j, k \in [1, \dots, L]$, $1 \leq i \leq n_0$.

Taking $i = 0$, we see that Condition A4(a) implies $\{T^{-1}(x)\} \neq \emptyset$ for every $x \in [1 - a, 1]$, i.e. the hole is not allowed to eliminate both preimages of any point. Given the small size of the holes, Condition A4 can be interpreted as requiring that the holes be in generic position with respect to one another.

2.3. Statement of results.

THEOREM 2.2. *Given a logistic map $\hat{T} \in \mathcal{M}$ and a hole H in $[-1, 1]$ satisfying Conditions A1–A3, the open system (T, I) has a Markov extension (F, Δ) with an exponential rate of return.*

Projecting the a.c.c.i.m. for (F, Δ) onto I yields the second theorem.

THEOREM 2.3. *Under the conditions of Theorem 2.2, T admits an absolutely continuous conditionally invariant measure on I . The density ψ can be written as*

$$\psi = \rho_1 + \rho_2$$

where ρ_1 is of bounded variation and

$$\rho_2(x) \leq \text{const.} \sum_{k=1}^{\infty} \frac{(1.9)^{-k/3}}{\sqrt{x - \hat{T}^k(0)}}.$$

If in addition Condition A4 is satisfied, then ψ is bounded away from zero on $[1 - a, 1] \setminus H$.

We obtain convergence of these measures as the size of the hole goes to zero.

THEOREM 2.4. *Let \hat{T} and H satisfy Conditions A1–A4. Define $H_t = H$ and let $\{H_s\}$ for $s \in [0, t]$ be a sequence of holes with the following properties:*

- (1) $mH_s \leq s$, $H_s \subset H_t$ and each component of H_t contains at most one component of H_s ;
- (2) either $0, \delta, -\delta \notin H_t$ or $0, \delta, -\delta \in H_s$ for all $s \in [0, t]$.

Let μ_s be the a.c.c.i.m. corresponding to H_s obtained from Theorem 2.3. As s goes to zero, the sequence μ_s converges weakly to the unique absolutely continuous invariant measure for \hat{T} with no holes.

The conditions on the sequence of holes ensure that the intervals of monotonicity of the map T do not decrease in length. This allows us to choose the same constants in our construction for each s and so gain uniform estimates.

2.4. Some properties of maps in the class \mathcal{M} . Before beginning our construction of the Markov extension of a logistic map \hat{T} , we review some properties of maps in the class \mathcal{M} defined in § 2.1.

The following lemma is proved in [WY2]. We present the proof here for completeness.

LEMMA 2.5. For $\hat{T} \in \mathcal{M}$, there exists $c'_0 > 0$ such that the following hold for all $\delta < \delta_0$:

- (a) if $x, \hat{T}(x), \dots, \hat{T}^{n-1}(x) \notin (-\delta, \delta)$, then $|(\hat{T}^n)'(x)| \geq c'_0 \delta e^{\frac{1}{3}\lambda_0 n}$;
- (b) if $x, \hat{T}(x), \dots, \hat{T}^{n-1}(x) \notin (-\delta, \delta)$ and $\hat{T}^n(x) \in (-\delta_0, \delta_0)$, then $|(\hat{T}^n)'(x)| \geq c_0 e^{\frac{1}{3}\lambda_0 n}$.

Proof. Let x satisfy $\hat{T}^i(x) \notin (-\delta, \delta)$ for $i = 1, \dots, n-1$. Suppose $\hat{T}^i(x)$ enters $(-\delta_0, \delta_0)$ at times t_1, \dots, t_s before time n . Let $t_0 = 0$ and $t_{s+1} = n$. We set $k_j = t_{j+1} - t_j$ and estimate the derivative on each time interval $[t_j, t_{j+1}]$. There are four cases to consider.

Case 1. $\hat{T}^{t_j}(x), \hat{T}^{t_{j+1}}(x) \in (-\delta_0, \delta_0)$. Then $|(\hat{T}^{k_j})'(\hat{T}^{t_j}x)| \geq e^{\frac{1}{3}\lambda_0 k_j}$ using Definition 2.1(b)(ii) and (c).

Case 2. $\hat{T}^{t_j}(x) \notin (-\delta_0, \delta_0), \hat{T}^{t_{j+1}}(x) \in (-\delta_0, \delta_0)$. Then $|(\hat{T}^{k_j})'(\hat{T}^{t_j}x)| \geq c_0 e^{\lambda_0 k_j}$ using Definition 2.1(b)(ii).

Case 3. $\hat{T}^{t_j}(x), \hat{T}^{t_{j+1}}(x) \notin (-\delta_0, \delta_0)$. If $k_j \geq M_0$, then $|(\hat{T}^{k_j})'(\hat{T}^{t_j}x)| \geq e^{\lambda_0 k_j}$ by Definition 2.1(b)(i). Otherwise, $|(\hat{T}^{k_j})'(\hat{T}^{t_j}x)| \geq c''_0 e^{\frac{1}{3}\lambda_0 k_j}$ with $c''_0 = \delta_0^{M_0-1} e^{-\frac{1}{3}\lambda_0(M_0-1)}$.

Case 4. $\hat{T}^{t_j}(x) \in (-\delta_0, \delta_0), \hat{T}^{t_{j+1}}(x) \notin (-\delta_0, \delta_0)$. This is the same as Case 3 with an additional factor $\geq \delta$.

Stringing these cases together, we obtain (b) using cases 1 and 2 and (a) with $c'_0 := c_0 \cdot c''_0$. \square

It may seem at first glance that the introduction of a new $\delta < \delta_0$ is redundant since there are analogous properties associated with each. The key difference, however, is that δ_0 and the constants associated with it depend only on the map \hat{T} , whereas we are free to choose δ . We shall choose δ depending on several factors involved in the construction of the tower as well as the placement of the hole.

We now explore a second property of maps in the class \mathcal{M} . This property concerns a period of recovery for $(\hat{T}^j)'(x)$ for orbits which pass through a δ -neighborhood of the critical point. Let $\delta = e^{-k_0}$ and define a partition of $(-\delta, \delta)$ into intervals $I_k = (e^{-(k+1)}, e^{-k})$, $k \geq k_0$, and $I_k = -I_{-k}$ for $k \leq -k_0$. k_0 will be chosen large enough so that the series in the proof of Proposition 4.1 converge.

For $x \in I_k$, define $\tilde{p}(x) = \max\{n \in \mathbb{Z} : |\hat{T}^j x - \hat{T}^j 0| < 1/j^2, \text{ for all } j < n\}$. Let

$$p(x) = \inf_{y \in I_k} \tilde{p}(y).$$

The number $p(x)$ is called the *bound period* of x by Benedicks and Carleson in [BC]. We call an interval $\omega \subset I_k$ *bound* from time 1 until time $p-1$ and *free* from time p until ω enters $(-\delta, \delta)$ again. Then another bound period begins. Since p is constant on each I_k , we sometimes refer to p as $p(k)$. For a near 2, the following properties of p are proved in [BC] and outlined succinctly in [Y1].

PROPERTIES P1. The function $p : (-\delta, \delta) \rightarrow \mathbb{Z}^+$ is constant on each I_k and increasing with $|k|$. In addition, for $x \in I_k$:

- (a) $\frac{1}{2}|k| \leq p(x) \leq 4|k|$;
- (b) $|(\hat{T}^j)'(x)| \approx \text{const.} |(\hat{T}^j)'(\hat{T}^j 0)| \geq \text{const.} (1.9)^j$, for all $j < p(x)$;
- (c) $|(\hat{T}^p)'(x)| \geq e^{p/5}$.

The central distortion estimate which yields Properties P1 is given at the beginning of §4.5.

We choose δ small enough so that I_{k_0} is free by the time it leaves an $(r/2)$ -neighborhood of the critical orbit. This in turn implies that any interval $\omega \subset I_k$ must be free at time n if $\hat{T}^n \omega$ intersects H . But we may conclude more than this. In fact, Condition A1 together with Lemma 4.5 ensures that *each I_k must grow to a fixed size before intersecting the hole*. We call this fixed length ε' .

We define time $q(k)$ for $k \geq k_0$ by

$$q(k) = \max\{n \in \mathbb{Z} : |\hat{T}^j(0, e^{-k})| \leq r/2, \forall j \leq n\}.$$

$q(k)$ is defined analogously for $k \leq -k_0$. Using Lemma 4.5, we shall prove in §4.5 that each I_k must grow to length ε' by time $q(k)$. We also use $q(k)$ to define our construction of the stopping time S and partition \mathcal{Z} of Proposition 4.1. Note that $q(k) \geq p(k)$.

2.5. Markov extensions. We describe the main ideas of the construction of a Markov extension for maps of the interval following [Y3]. We carry out this construction in detail in §4.

Given a subinterval Λ and a map T , we consider the forward images of Λ under the action of T . When a connected component of $T^n \Lambda$ covers Λ , we declare ω , the subinterval of Λ satisfying $T^n \omega = \Lambda$, to have returned and stop iterating it. We continue to iterate the remaining components of $T^n \Lambda$ until they return to completely cover Λ . In this way, we generate a countable partition $\{\Lambda_i\}$ of subintervals of Λ and a stopping time $R : \Lambda \rightarrow \mathbb{N}$, constant on elements of the partition and satisfying $T^R(\Lambda_i) = \Lambda$. Then $\{\Lambda_i\}$ is a countable Markov partition for the map T^R .

In this situation, we define a *Markov extension* of $T : \bigcup_{n \geq 0} T^n \Lambda \hookrightarrow$ as a dynamical system $F : \Delta \hookrightarrow$ for which there exists a projection $\pi : \Delta \rightarrow \bigcup_{n \geq 0} T^n \Lambda$ such that $\pi \circ F = T \circ \pi$.

We also call $F : \Delta \hookrightarrow$ the *tower model* or simply the tower associated with T . The reason for this is the following pictorial model for the Markov extension. Let $\Delta_0 = \Lambda$ and define

$$\Delta = \{(x, n) \in \Delta_0 \times \mathbb{N} : R(x) > n\}.$$

The tower map is given by

$$F = \begin{cases} F(x, n) = (x, n+1) & \text{if } n+1 < R(x), \\ F(x, n) = (T^R(x), 0) & \text{if } n+1 = R(x). \end{cases}$$

The l th level of the tower is $\Delta|_{n=l}$ and the action of the tower map F is to map a point up the levels of the tower until time R at which time the point is returned to the base Δ_0 . Note that all of the returns to the base are Markov because of the nature of the returns of Λ_i to Λ .

The flexibility of the Markov extension stems from the fact that the dynamical system in question need not be uniformly hyperbolic. What matters is the average behavior of the map T between returns to Λ . This is what allows the method to be applied to Hénon maps and the logistic family. There are three basic steps which are required for this method to work:

- (1) given a dynamical system $T : M \curvearrowright$, we construct a Markov extension $F : \Delta \curvearrowright$;
- (2) we prove results about (F, Δ) using its simpler properties: namely, controlled hyperbolicity and a countable Markov structure with a certain decay rate in the measure of the elements of the partition;
- (3) we pass these results back to the original system (T, M) .

Step (1) is completed by the construction contained in §4. This is the most technical part of the paper. Step (2) is proved in [D] and those results are recalled in §3. Step (3) is completed in §5.

3. Tower maps with holes

The results of [D] for tower maps with holes apply in a more general setting than the present paper. Logistic maps are C^2 and the tower which we construct will have no holes in its base. Here we recall only those results relevant to our case. This simplifies the assumptions on the tower somewhat. We do, however, retain the definition of the function space X in which the conditionally invariant density lies since we will use this to establish the properties of the conditionally invariant density for the logistic map in §5.

3.1. Tower with multiple bases. The towers studied in [D] are towers with multiple bases $\hat{\Delta}_0^{(i)}$, which are intervals of unit length whose interiors are pairwise disjoint. The base $\hat{\Delta}_0 = \bigcup_{i=1}^N \hat{\Delta}_0^{(i)}$ is also an interval. We let m denote one-dimensional Lebesgue measure on the tower and let \mathcal{Z} be a countable partition of $\hat{\Delta}_0$ whose elements are subintervals of the $\hat{\Delta}_0^{(i)}$. Given a return time function $R : \hat{\Delta}_0 \rightarrow \mathbb{Z}^+$ which is constant on each element of \mathcal{Z} , a tower $(\hat{\Delta}, \hat{F}, m)$ is built over $\hat{\Delta}_0$ with

$$\hat{\Delta} := \{(z, n) \in \hat{\Delta}_0 \times \mathbb{N} : n < R(z)\}.$$

As before, we call the l th level of the tower $\hat{\Delta}_l := \hat{\Delta}|_{n=l}$ and $\hat{\Delta}_l^{(i)}$ is the part of $\hat{\Delta}_l$ directly over $\hat{\Delta}_0^{(i)}$. We let $\hat{\Delta}^{(i)} = \bigcup_{l=0}^{\infty} \hat{\Delta}_l^{(i)}$.

The action of $\hat{F} : \hat{\Delta} \rightarrow \hat{\Delta}$ is $\hat{F}(z, l) = (z, l+1)$ if $l+1 < R(z)$ and $\hat{F}^{R(z)}(\mathcal{Z}(z)) = \hat{\Delta}_0^{(i)}$ for some $1 \leq i \leq N$, where $\hat{F}^{R(z)}|_{\mathcal{Z}(z)}$ is continuous and one-to-one and $\mathcal{Z}(z)$ is the element of \mathcal{Z} containing z .

The first assumption made on the tower is that the measure of the levels of the tower decays exponentially. This is crucial to the existence of an a.c.c.i.m. with good properties.

Condition H1. There exist $A > 0$ and $0 < \theta < 1$ such that $m(\hat{\Delta}_l) \leq A\theta^l$ for $l \geq 0$.

We leave assumptions about the regularity of \hat{F} until after we have introduced the holes.

3.2. Introduction of holes and regularity of \hat{F} . A hole \tilde{H} in $\hat{\Delta}$ is a union of open intervals $\tilde{H}_{l,j}^{(i)}$ such that $\bigcup_j \tilde{H}_{l,j}^{(i)} =: \tilde{H}_l^{(i)} \subset \hat{\Delta}_l^{(i)}$ with finitely many $\tilde{H}_{l,j}^{(i)}$ per level l . We set $\tilde{H}_l = \bigcup_{i=1}^N \tilde{H}_l^{(i)}$. We require that each $\tilde{H}_{l,j}^{(i)} = \hat{F}^l(\omega)$, where ω is the union of elements of \mathcal{Z} , thus preserving the Markov structure of the returns to $\hat{\Delta}_0$. If $\hat{F}^l(\omega) = \tilde{H}_{l,j}^{(i)}$, then the intervals on all levels of the tower directly above $\tilde{H}_{l,j}^{(i)}$ are deleted since once \hat{F} maps a point into \tilde{H} , it disappears forever. ω does not return to $\hat{\Delta}_0$.

Let $\Delta = \hat{\Delta} \setminus \tilde{H}$ and $\Delta_l = \hat{\Delta}_l \setminus \tilde{H}$ with analogous definitions for $\Delta^{(i)}$ and $\Delta_l^{(i)}$. We assume the existence of a countable Markov partition $\{\Delta_{l,j}^{(i)}\}$ with $\bigcup_j \Delta_{l,j}^{(i)} = \Delta_l^{(i)}$ for each i and l . Each $\Delta_{l,j}^{(i)}$ is an interval comprised of countably many elements of the form $\hat{F}^l(\omega)$, $\omega \in \mathcal{Z}$, and $\hat{F}|_{\Delta_{l,j}^{(i)}}$ is one-to-one.

In applications, $\{\Delta_{l,j}^{(i)}\}$ is dynamically defined during the construction of the tower and its elements are the maximal intervals which project onto the iterated pieces of the reference set Δ at time l . For this reason, it is useful to keep track of the elements $\Delta_{l,j}^{(i)}$ rather than the elements of $\hat{F}^l(\mathcal{Z})$.

We denote by $\Delta_{l,j}^{(i)*}$ those $\Delta_{l,j}^{(i)}$ whose image returns to the base, i.e. such that $\hat{F}(\Delta_{l,j}^{(i)}) = \bigcup_{k=k_1}^{k_2} \Delta_0^{(k)}$ for some $1 \leq k_1 \leq k_2 \leq N$, and set $\Delta^* = \bigcup \Delta_{l,j}^{(i)*}$.

PROPERTIES P2. Since \hat{T} is C^2 , the map \hat{F} has the following properties with respect to the partition $\{\Delta_{l,j}^{(i)}\}$.

- (a) \hat{F} is C^2 on each $\Delta_{l,j}^{(i)}$.
- (b) There exist $\gamma > 1$ and $\beta > 0$ such that on $\Delta_{l,j}^{(i)*}$, $|\hat{F}'| \geq \gamma e^{\beta l}$. Elsewhere, $|\hat{F}'| = 1$.
- (c) *Bounded Distortion.* There exists $C > 0$ such that for any $x, y \in \hat{\Delta}_0^{(i)}$ and $x', y' \in \Delta_{l,j}^{(i)*}$ such that $\hat{F}(x') = x$ and $\hat{F}(y') = y$ we have

$$\left| \frac{\hat{F}'(x')}{\hat{F}'(y')} - 1 \right| \leq C|x - y|. \quad (2)$$

Controlling the distortion for logistic maps requires a countable partition in a neighborhood of the critical point. Such a partition has been introduced in §2.4 and equation (2) is a consequence of the distortion lemmas of §4.5.

Let $F = \hat{F}|_{(\Delta \setminus \hat{F}^{-1}\tilde{H})}$. We say F is *transitive on components* if for all pairs i, j , $1 \leq i, j \leq N$, there exists an m such that $F^m \Delta_0^{(i)} \supseteq \Delta_0^{(j)}$. Note that if $N = 1$, then transitivity on components is automatic as long as the hole allows at least one return to the base.

3.3. *Definition of a convex functional.* The Perron–Frobenius operator associated with F acts on $L^1(\Delta)$ by

$$\mathcal{P}f(x) = \sum_{y \in F^{-1}x} \frac{f(y)}{|F'(y)|}.$$

We define $\mathcal{P}_1 f = \mathcal{P}f / \|\mathcal{P}f\|_{L^1}$ and seek a fixed point for the operator \mathcal{P}_1 . A fixed point for \mathcal{P}_1 is a conditionally invariant density for F .

Choose $\xi > 0$ small enough so that $e^{-\xi} > \max\{\theta, e^{-\beta}\}$. Given $f \in L^1(\Delta)$, let $f_{l,j}^{(i)} = f|_{\Delta_{l,j}^{(i)}}$. Let $\|f\|_\infty$ denote the L^∞ norm of f and define

$$\|f_{l,j}^{(i)}\|_\infty = \|f_{l,j}^{(i)}\|_\infty e^{-\xi l}, \quad \|f_{l,j}^{(i)}\|_r = \sup_{\substack{x \in \Delta_{l,j}^{(i)} \\ f(x) \neq 0}} \left| \frac{f'(x)}{f(x)} \right| e^{-\xi l}.$$

Then define

$$\|f\| = \max\{\|f\|_\infty, \|f\|_r\},$$

where $\|f\|_\infty = \sup_{i,l,j} \|f_{l,j}^{(i)}\|_\infty$ and $\|f\|_r = \sup_{i,l,j} \|f_{l,j}^{(i)}\|_r$. Let $X = \{f : \Delta \rightarrow \mathbb{C} \mid \|f\| < \infty\}$. Although $\|\cdot\|_r$ is not a norm, it does satisfy a convex-like inequality on a subset X_M of X defined by

$$X_M = \left\{ f \in X \mid \|f\| \leq M, f \geq 0, \int_\Delta f \, dm = 1 \right\}.$$

It is proved in [D] that X_M is a convex, compact subset of $L^1(\Delta)$. We take $M = b/(1-a_0)$, where a_0 and b are defined below.

3.4. Condition on the holes and main result. We formulate a single condition involving the measure of the holes which guarantees the existence of an a.c.c.i.m. in X .

Let $a_0 := \max\{e^{-\xi}, 1/\gamma\}$ and $b := 1 + C$. The required condition on the holes is as follows:

Condition H2.

$$\sum_{l \geq 1} e^{\xi(l-1)} m \tilde{H}_l \leq \frac{(1-a_0)^2}{b}.$$

The main result we wish to recall from [D] is the following theorem. We will apply this theorem in §5 after constructing the Markov extension.

THEOREM 3.1. *Given a tower with holes (Δ, F, m) with Properties P2 and under Conditions H1 and H2, there exists a probability density φ in X_M such that $\mathcal{P}_1 \varphi = \varphi$. If in addition F is transitive on components, then φ is the unique non-trivial conditionally invariant density in X and φ is bounded away from zero on Δ .*

Remark. Note that since $\varphi \in X$, its eigenvalue λ must satisfy $\lambda \geq e^{-\xi}$. In fact, in [D] it is proven that $\lambda \geq 1 - M \sum_{l \geq 1} e^{\xi(l-1)} m \tilde{H}_l$. This estimate stems from the lower bound on the renormalization constant for functions in the set X_M .

4. Construction of the tower

In this section we describe the construction of the Markov extension of $T : I^1 \rightarrow I$. The construction entails finding the right length scale for the reference intervals $\Lambda^{(i)}$ which will constitute the base of the tower and showing that the object we construct has certain properties. These are summarized in Proposition 4.2 in §4.3 and Proposition 4.3 in §4.4.

The construction of the tower involves a series of constants which we define below. Some have been introduced already. The order of their selection is important and follows that of the list.

- The constants δ_0 and λ_0 introduced in Definition 2.1. Throughout the proofs of §4, for definiteness we consider λ_0 as $\log 1.9$.
- The minimum distance r between H and the critical orbit introduced by Condition A1.
- m_0 (depending on λ_0) and ε_0 introduced in Condition A2. If λ_0 is taken to be $\log 1.9$, then $m_0 = 10$ is large enough.

- n_0 (depending on ϵ_0) is the least i for which $\hat{T}^i J \supseteq [1 - a, 1]$ for every interval J of length at least $\epsilon_0/2$. n_0 is used in Condition A4 and later in §5.3 to prove a transitivity property for the map with holes.
- $\delta = e^{-k_0}$, which defines a δ -neighborhood of the critical point and induces the partition $\{I_k\}_{|k| \geq k_0}$ defined in §2.4. δ is chosen small enough to make the series in the proof of Proposition 4.1 converge and also so that I_{k_0} is free by the time it leaves an $(r/2)$ -neighborhood of the critical orbit.
- ϵ' , the fixed length to which every I_k must grow by time $q(k)$ before intersecting the hole, proven after Lemma 4.5.
- ϵ , the length of the reference intervals $\Lambda^{(i)}$ which constitute the base of the tower. ϵ is chosen so that $4^8 \epsilon = \min\{\epsilon', \epsilon_0, 1/4\tilde{C}\}$ where \tilde{C} is the nonlinearity constant in the distortion estimate of Lemma 4.6. Since $\tilde{C} \sim 1/\delta^2$, this requires $\epsilon \sim \delta^2$. ϵ is chosen to be small compared to ϵ' and ϵ_0 in order to control the rate at which pieces are generated during the construction of the tower. The requirement involving \tilde{C} ensures a minimum expansion at the return time.

We begin by defining a partition \mathcal{Q} and a type of interval Ω which we shall use in our construction.

Recall the partition of $(-\delta, \delta)$ introduced earlier: $\{I_k\}_{|k| \geq k_0}$. To this partition we join the partition of $[-1, 1]$ into the finitely many maximal intervals of I and H . We call this new partition \mathcal{Q} .

Let Ω be an interval such that $\epsilon \leq |\Omega| \leq 3\epsilon$. We require that $\Omega \subset I$ and that either $\Omega \subset (-\delta, \delta) \setminus \{0\}$ or $\Omega \subset I \setminus (-\delta, \delta)$.

We cover $[1 - a, 1] \setminus H$ with intervals $\Lambda^{(1)}, \dots, \Lambda^{(N)}$, each of which is of the type Ω described above, except that we restrict $\epsilon \leq |\Lambda^{(i)}| \leq 2\epsilon$. The intervals $\Lambda^{(i)}$ are the reference intervals which will serve as the base of the tower.

4.1. Introduction of an auxiliary stopping time. Let Ω be an interval of the form described above. The principal properties of the auxiliary stopping time and partition that we shall construct on Ω are listed in the following proposition.

PROPOSITION 4.1. *There exist a countable partition \mathcal{Z} of Ω and a stopping time S satisfying:*

- S is constant on each element $\omega \in \mathcal{Z}$;
- $T^S \omega$ is defined and $|T^S \omega| \geq 4^8 \epsilon$ or $T^{S-1} \omega$ is defined and $\hat{T}^S \omega \subset H$;
- There exists $\tilde{C} \sim 1/\delta^2$ such that for $x, y \in \omega$,

$$\left| \frac{(\hat{T}^S)'(x)}{(\hat{T}^S)'(y)} - 1 \right| \leq \tilde{C} |\hat{T}^S(x) - \hat{T}^S(y)|;$$

- $|(\hat{T}^S)'(x)| > 4^6$;
- $m\{x \in \Omega : S(x) > n\} \leq C' e^{-n/21}$ for some C' independent of δ .

The proof of this proposition in §§4.1 and 4.2 follows closely the approach of Benedicks and Young [BY] for Hénon maps without holes.

We construct \mathcal{Z} and S as follows. We take components of $\mathcal{Q} \setminus \Omega$ and place them in the set Ω_0 . Given $\Omega_{n-1} \subset \Omega$, we proceed inductively. Let $\omega \in \Omega_{n-1}$. Let t be the last time ω

passed through $(-\delta, \delta)$ and let k be such that $\hat{T}^t \omega \subset I_k$. If ω has not yet passed through $(-\delta, \delta)$ by time n , set $t = q(k) = 0$.

If $n > t + q(k)$, we look at $\hat{T}^n \omega$ and do the following.

Case 1. $\hat{T}^n \omega$ does not intersect the hole. If $|\hat{T}^n \omega| \geq 4^8 \varepsilon$, then enter ω as an element of \mathcal{Z} and declare the stopping time $S(x) \equiv n$ on ω . Otherwise partition ω according to $\hat{T}^{-n} \mathcal{Q}|\omega$ and put these pieces into Ω_n . If $\hat{T}^n \omega$ lies partly outside and partly inside of $(-\delta, \delta)$ then we append the piece lying outside to the piece of $\hat{T}^n \omega$ lying in $I_{\pm k_0}$ and do not introduce a cut there. Since $\varepsilon \sim \delta^2$, this added length is negligible from time n to time $n + q(k_0)$.

Case 2. $\hat{T}^n \omega$ intersects the hole. Set $S(x) \equiv n$ on the components of $\omega \cap \hat{T}^{-n} H$ and enter them as elements of \mathcal{Z} . Take the remaining subintervals of ω and follow the procedure described in Case 1 for each. Note that there can be at most two subintervals ω_n of ω such that $|\hat{T}^n \omega_n| < 4^8 \varepsilon$ because of Condition A2 and our choosing $4^8 \varepsilon \leq \varepsilon_0$.

If $n < t + q(k)$, then $\mathcal{Q}|\hat{T}^n \omega$ will have only one component. We put ω in Ω_n and continue to iterate it.

If $n = t + q(k)$, then if $|\hat{T}^n \omega| < 4^8 \varepsilon$, we put $\omega \in \Omega_n$. If $|\hat{T}^n \omega| \geq 4^8 \varepsilon$, we do one of two things.

Case 1. $\omega \in \Omega_{t-1}$. Then ω was not created by a cut at time t . We declare $S(x) \equiv n$ on ω and enter ω as an element of \mathcal{Z} .

Case 2. $\omega \notin \Omega_{t-1}$. Then ω was created at time t by a cut between I_k and I_{k+1} . So there are two intervals ω and γ such that $\omega \cup \gamma$ is one interval until time t , $\hat{T}^t \omega \subseteq I_k$ and $\hat{T}^t \gamma \subseteq I_{k+1}$; but $\hat{T}^n \omega$ and $\hat{T}^n \gamma$ are still adjacent. $\hat{T}^n \omega$ will overlap a large number of the $\Lambda^{(i)}$. On the side of ω adjacent to γ , we adjoin to γ the part of ω which does not completely cover the last $\Lambda^{(i)}$ on that side under \hat{T}^n . Let us call this interval ω' . We declare $S(x) \equiv n$ on $\omega \setminus \omega'$ and put the interval $\omega' \cup \gamma$ into Ω_n and continue to iterate it. We do this to control the number of pieces generated by the process described later in §4.3. We will need this control in order to obtain the bounds on the conditionally invariant density in §5.4. (Note that if ω had been created by a cut between I_k and I_{k-1} , the process of adjoining a left over piece on that side would already have occurred at time $t + q(k - 1)$.)

It is clear that Proposition 4.1(a) and (b) will be satisfied by the construction described above. Item (c) is proven by the distortion bounds of Lemma 4.6 and item (d) will follow immediately from that. Item (e) is proved in §4.2.

We close this section by showing that every interval of length at least ε will grow to length $4^8 \varepsilon$ using the upper bound on the size of the hole given by equation (1). Suppose Ω is an interval of length at least ε and suppose that Ω intersects the hole after its very first iterate. Then there will be at most two pieces of Ω whose image did not fall into the hole. Choose the longer of the two pieces and call it ω_1 . Using Lemma 2.5(a), we observe that

$$|\omega_1| \geq \frac{1}{2} \left(\varepsilon - \frac{mH}{c'_0 \delta e^{\frac{1}{3}\lambda_0}} \right).$$

If ω_1 does not grow to length $4^8 \varepsilon$, it must wait at least another m_0 iterates before intersecting H again. Say this happens at time t_1 . Once again, there are at least two pieces of ω_1 whose images do not intersect the hole under \hat{T}^{t_1} . Call the longer of these ω_2

and note that

$$|\omega_2| \geq \frac{1}{2} \left(\frac{1}{2} \left(\varepsilon - \frac{mH}{c'_0 \delta e^{\frac{1}{3}\lambda_0}} \right) - \frac{mH}{c'_0 \delta e^{\frac{1}{3}\lambda_0(m_0+1)}} \right).$$

Repeating this process k times and always following the larger half, we see that

$$|\omega_k| \geq \frac{\varepsilon}{2^k} - \sum_{i=0}^{k-1} \frac{mH}{c'_0 \delta e^{\frac{1}{3}\lambda_0(im_0+1)} 2^{k-i}} \geq \frac{\varepsilon}{2^{k+1}},$$

where we have used equation (1) in the last step. Following this process until time n , and noting that $n \geq m_0 k$, we have

$$|\hat{T}^n \omega_k| \geq c'_0 \delta e^{\frac{1}{3}\lambda_0 n} \frac{\varepsilon}{2^{(n/m_0)+1}},$$

which is exponentially increasing. This will continue until a part of Ω grows to length $4^8 \varepsilon$. If along the way, $\hat{T}^n \omega_k$ lands in $(-\delta, \delta)$, then our estimates only improve since the piece cannot intersect the hole again until the partition element it lies in grows to size $\varepsilon' \geq 4^8 \varepsilon$.

4.2. Estimating the return time function S . In this section we prove that $m\{x \in \Omega : S(x) > n\} \leq C' e^{-n/2^1}$, which is part (e) of Proposition 4.1.

In order to estimate the tail of the return time function S , we will use information about the times when an interval passes through $(-\delta, \delta)$. Recall that for $\omega \in \Omega_{n-1}$ if $\hat{T}^n \omega$ intersects $(-\delta, \delta)$, then we introduce cuts in ω according to the partition $\hat{T}^{-n} \mathcal{Q}|\omega$ and the pieces are entered as elements of Ω_n . We keep track of which interval I_k each piece passes through at time n .

If an interval ω is a subset of I_{r_i} at time t_i , $1 \leq i \leq s$, then we say ω has *itinerary* (r_1, \dots, r_s) . Let $p_i = p(I_{r_i})$. (p_1, \dots, p_s) are the recovery times associated with the itinerary (r_1, \dots, r_s) . Recall that if $\hat{T}^n \omega$ lies partly outside of $(-\delta, \delta)$ then we append the piece lying outside to the piece of $\hat{T}^n \omega$ lying in $I_{\pm k_0}$ and do not introduce a cut there. This will not affect the recovery time $p(\pm k_0)$ of $I_{\pm k_0}$.

Notice that by construction, pieces that are created by an interval landing on $(-\delta, \delta)$ at any given time will have different itineraries; however, if an interval is mapped across one of the holes H_j and split into two pieces, then those two pieces may be mapped into $(-\delta, \delta)$ at different times and so generate separate pieces with the same itinerary. We wish to obtain an upper bound on the number of pieces with the same itinerary up to time n that can be created from a single interval which is iterated according to the procedure described after the statement of Proposition 4.1.

Let $\omega \subset I_{r_0}$ and let S_n be the set of elements of Ω_n which have the same itinerary (r_1, \dots, r_s) at time n . Now I_{r_0} cannot intersect the hole (and generate more pieces) for the first p_0 iterates. Then from time p_0 to time t_1 , it can be cut at most $1 + [(t_1 - p_0)/m_0]$ times, where $[\cdot]$ denotes the greatest integer function. This will be true on each

time interval $[t_i, t_{i+1}]$. Thus

$$\begin{aligned} \log_2(\#S_n) &\leq 1 + \left\lfloor \frac{t_1 - p_0}{m_0} \right\rfloor + 1 + \left\lfloor \frac{t_2 - (t_1 + p_1)}{m_0} \right\rfloor + \dots \\ &\quad + 1 + \left\lfloor \frac{t_s - (t_{s-1} + p_{s-1})}{m_0} \right\rfloor + 1 + \left\lfloor \frac{n - (t_s + p_s)}{m_0} \right\rfloor \\ &\leq \begin{cases} s + 1 + \frac{n - \sum_{i=0}^s p_i}{m_0} & \text{if } n \geq t_s + p_s \\ s + \frac{t_s - \sum_{i=0}^{s-1} p_i}{m_0} & \text{if } n < t_s + p_s \end{cases} \end{aligned} \quad (3)$$

$$\leq s + 1 + \frac{n - \frac{1}{2} \sum_{i=0}^{s-1} |r_i|}{m_0}, \quad (4)$$

where we have used Property P1(a) in the last step. Now we are ready to estimate the tail of the return time function S .

We begin with an interval Ω which may or may not be a subset of $(-\delta, \delta)$. Suppose $\omega \in \Omega_n$ has itinerary (r_0, \dots, r_s) at times t_0, \dots, t_s with $s \geq 1$ and $t_0 = 0$. If $\omega \subset I_k$, let $r_0 = k$; otherwise, let $r_0 = 0$. Assume $|k| \leq n/8$. For $x \in \omega$,

$$|(\hat{T}^n)'(x)| = |(\hat{T}^{n-t_s})'(\hat{T}^{t_s}x)| \prod_{i=0}^{s-1} |(\hat{T}^{t_{i+1}-t_i})'(\hat{T}^{t_i}x)|.$$

We estimate $|(\hat{T}^{t_{i+1}-t_i})'(\hat{T}^{t_i}x)|$ using a method similar to the proof of Lemma 2.5. Let $s_0 = t_i$, $s_{k+1} = t_{i+1}$ and s_1, \dots, s_k be the times when ω returns to $(-\delta_0, \delta_0)$ between times t_i and t_{i+1} . On each interval $[s_j, s_{j+1}]$ (except possibly when $i = 0$ and $j = 0$), we are in Case 1 of the proof of Lemma 2.5 so $|(\hat{T}^{s_{j+1}-s_j})'(\hat{T}^{t_i+s_j}x)| \geq e^{\frac{1}{3}\lambda_0(s_{j+1}-s_j)}$. Stringing these intervals together, we have

$$|(\hat{T}^{t_{i+1}-t_i})'(\hat{T}^{t_i}x)| \geq e^{\frac{1}{3}\lambda_0(t_{i+1}-t_i)} \quad (5)$$

for $i > 0$ and $|(\hat{T}^{t_1})'(x)| \geq c_0 e^{\frac{1}{3}\lambda_0 t_1}$. This yields

$$|(\hat{T}^n)'(x)| \geq |(\hat{T}^{n-t_s})'(\hat{T}^{t_s}x)| \cdot c_0 e^{\frac{1}{3}\lambda_0 t_s}.$$

If $n \geq t_s + p_s$, then $|(\hat{T}^{n-t_s})'(\hat{T}^{t_s}x)| \geq c'_0 \delta e^{\frac{1}{3}\lambda_0(n-t_s-p_s)} e^{p_s/5}$ using Property P1(c) and Lemma 2.5(a). Combining these estimates, we have

$$|(\hat{T}^n)'(x)| \geq c_0 \cdot c'_0 \delta e^{\frac{1}{3}\lambda_0 n}. \quad (6)$$

Since $|\hat{T}^n \omega| < 4^8 \varepsilon$, we can estimate

$$|\omega| \leq \frac{4^8 \varepsilon}{\delta c_0 c'_0 e^{\frac{1}{3}\lambda_0 n}} \leq \frac{1}{c_0 c'_0} e^{-\frac{1}{3}\lambda_0 n}. \quad (7)$$

If $n < t_s + p_s$ then we note that at time t_s , $|(\hat{T}^{t_s})'(x)| \geq e^{\frac{1}{3}\lambda_0 t_s}$. Since $T^{t_s} \omega \subset I_{r_s}$, we estimate

$$|\omega| \leq \frac{e^{-|r_s|}}{c_0 e^{\frac{1}{3}\lambda_0 t_s}} \leq \frac{1}{c_0} e^{-\frac{1}{3}\lambda_0(t_s+4|r_s|)} e^{-|r_s|/8} \leq \frac{1}{c_0} e^{-\frac{1}{3}\lambda_0 n} e^{-|r_s|/8}. \quad (8)$$

We define $A(r_1, \dots, r_s) = \{x \in \omega : S(x) > n, x \text{ has itinerary } r_1, \dots, r_s\}$ and $K = \sum_{i=1}^s |r_i|$. For fixed s and K , we set $A_{s,K}^n = \bigcup_{\substack{(r_1, \dots, r_s) \\ \sum |r_i| = K}} A(r_1, \dots, r_s)$ and estimate

$$\#\{s\text{-tuples with } \sum |r_i| = K\} < 2^s \binom{K-1}{s-1}.$$

Note that since each $|r_i| \geq \Delta := \log 1/\delta$, we have $s \leq K/\Delta$. So

$$\#\{s\text{-tuples with } \sum |r_i| = K\} < 2^s \binom{K-1}{K/\Delta-1} < 2^s (1 + \sigma(\delta))^K,$$

where $\sigma(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. Now we estimate

$$m(A_{s,K}^n) \leq \left(\#\{s\text{-tuples with } \sum |r_i| = K\} \right) \cdot \left(\begin{array}{c} \# \text{ pieces with} \\ \text{itinerary } (r_1, \dots, r_s) \end{array} \right) \cdot m \left\{ \begin{array}{c} \text{one piece with} \\ \text{itinerary } (r_1, \dots, r_s) \end{array} \right\}. \quad (9)$$

Let Ω' be those points in $\Omega \cap I_k$ with $S(x) > n$ which have made at least one return to $(-\delta, \delta)$.

$$m(\Omega') \leq \sum_{K \leq \frac{3}{2}n} \sum_{s=1}^{K/\Delta} m(A_{s,K}^n) + \sum_{K > \frac{3}{2}n} \sum_{s=1}^{K/\Delta} m(A_{s,K}^n). \quad (10)$$

For the first term, we use equation (4), the maximum of (7) and (8) and equation (9) to observe that

$$m(A_{s,K}^n) \leq 2^s (1 + \sigma(\delta))^K \cdot 2^{s+1} 2^{n/m_0} \cdot \frac{1}{c_0 c'_0} e^{-\frac{1}{3}\lambda_0 n}. \quad (11)$$

The competing factors in this expression are $e^{-\frac{1}{3}\lambda_0 n}$ and $2^{n/m_0}$. Taking $\lambda_0 = \log 1.9$ and $m_0 \geq 10$, we estimate their product by $e^{n((\log 2/m_0) - (\lambda_0/3))} \leq e^{-\frac{1}{7}n}$.

Since $n \geq \frac{2}{3}K$ and $n \geq 8|k| = 8|r_0|$, we may write $n \geq n/3 + K/3 + 8|r_0|/6$. Equation (11) becomes

$$m(A_{s,K}^n) \leq 2(1 + \sigma(\delta))^K 4^s \frac{1}{c_0 c'_0} e^{-\frac{1}{21}n} e^{-\frac{1}{21}K} e^{-\frac{4}{21}|r_0|}. \quad (12)$$

To estimate the second term of equation (10), we note that $t_s - \sum_{i=0}^{s-1} p_i \leq n - \frac{1}{2} \sum_{i=0}^{s-1} |r_i| = n - K/2 + |r_s|/2$. Applying this observation to equation (4) and using equation (8) we estimate

$$m(A_{s,K}^n) \leq 2^s (1 + \sigma(\delta))^K 2^{s+1} 2^{(n-K/2)/m_0} 2^{|r_s|/2m_0} \frac{1}{c_0} e^{-\frac{1}{3}\lambda_0(t_s+4|r_s|)} e^{-|r_s|/8}. \quad (13)$$

The competing factors in this expression are $e^{-\frac{1}{3}\lambda_0(t_s+4|r_s|)}$ and $2^{(n-K/2)/m_0}$. Using the fact that $n \leq \frac{2}{3}K$ and $t_s \geq \frac{1}{2} \sum_{i=0}^{s-1} |r_i|$, these terms become $e^{-\frac{1}{3}\lambda_0(K/2)} e^{-\frac{1}{3}\lambda_0(|r_0|/2)}$ and $2^{K/6m_0}$. Then if $m_0 \geq 10$ we conclude

$$m(A_{s,K}^n) \leq 2(1 + \sigma(\delta))^K 4^s \frac{1}{c_0} e^{-K/11} e^{-\frac{1}{6}\lambda_0|r_0|}. \quad (14)$$

Substituting equations (12) and (14) into equation (10) and summing over s , we get

$$\begin{aligned} m(\Omega') &\leq \sum_{K \leq \frac{3}{2}n} 2(1 + \sigma(\delta))^K 4^{(K/\Delta)+1} \frac{1}{c_0 c'_0} e^{-\frac{1}{21}K} e^{-\frac{1}{21}n} e^{-\frac{4}{21}|r_0|} \\ &\quad + \sum_{K > \frac{3}{2}n} 2(1 + \sigma(\delta))^K 4^{(K/\Delta)+1} \frac{1}{c_0} e^{-K/11} e^{-\frac{1}{6}\lambda_0|r_0|} \\ &\leq c_4 e^{-\frac{1}{21}n} e^{-\frac{4}{21}|r_0|} + \sum_{K > \frac{3}{2}n} c_5 e^{-K/30} e^{-\frac{1}{6}\lambda_0|r_0|} \\ &\leq c_6 e^{-n/21} e^{-\frac{1}{6}\lambda_0|r_0|} \end{aligned}$$

if δ is taken to be sufficiently small. Note that $\Delta > 50$ is forced by these estimates.

Let Ω'' be the set of points in $\Omega \cap I_k$ which have $S(x) > n$ and which have never returned to $(-\delta, \delta)$. In this case, we have a simple estimate using Lemma 2.5(a) that $|(\hat{T}^n)'(x)| \geq c'_0 \delta e^{\frac{1}{3}\lambda_0 n}$. Since we are assuming that $|k| \leq n/8$, we have $p_0 \leq 4|k| \leq n/2$ so using equation (4), the estimate on the number of pieces which can be formed up to time n , we have

$$m(\Omega'') \leq 2^{(n-p_0)/m_0} \frac{4^8 \varepsilon}{c'_0 \delta} e^{-\frac{1}{3}\lambda_0 n} \leq \frac{1}{c'_0} 2^{n/m_0} e^{-\frac{1}{3}\lambda_0 \frac{7}{8}n} e^{-\frac{1}{3}\lambda_0 |k|} \leq \frac{1}{c'_0} e^{-n/9} e^{-\frac{1}{3}\lambda_0 |r_0|}.$$

Now if Ω is not a subset of $(-\delta, \delta)$, then $r_0 = 0$ and we have shown that

$$m\{x \in \Omega : S(x) > n\} = m(\Omega') + m(\Omega'') \leq \left(c_6 + \frac{1}{c'_0}\right) e^{-n/21}.$$

On the other hand, if $\Omega \subset (-\delta, \delta)$, then

$$\begin{aligned} m\{x \in \Omega : S(x) > n\} &\leq \sum_{|k| \leq n/8} \left(c_6 + \frac{1}{c'_0}\right) e^{-n/21} e^{-\frac{1}{6}\lambda_0 |k|} + m(\Omega \cap (-e^{-n/8}, e^{-n/8})) \\ &\leq \left(c_6 + \frac{1}{c'_0}\right) e^{-n/21} \sum_{k \geq 1} e^{-\frac{1}{6}\lambda_0 k} + e^{-n/8} \\ &\leq c_7 e^{-n/21} \end{aligned}$$

which proves Proposition 4.1(e) with $C' = \max\{c_7, c_6 + 1/c'_0\}$.

4.3. Assembling the complete tower. We now have the required tools to complete the construction of the tower. This construction is achieved by applying Proposition 4.1 repeatedly to the reference intervals $\Lambda^{(i)}$ introduced at the beginning of §4. We fix j and proceed one $\Lambda^{(j)}$ at a time. Our construction will result in a partition and stopping time with the following properties.

PROPOSITION 4.2. *There exists a countable partition η of $\Lambda^{(j)}$ and a stopping time R satisfying:*

- (a) R is constant on each element $\omega \in \eta$;
- (b) either $T^R \omega$ is defined and $T^R \omega = \Lambda^{(i)}$ for some i , or $T^{R-1} \omega$ is defined and $\hat{T}^R \omega \subset H$;

(c) for $x, y \in \omega$,

$$\left| \frac{(\hat{T}^R)'(x)}{(\hat{T}^R)'(y)} - 1 \right| \leq \tilde{C} |\hat{T}^R(x) - \hat{T}^R(y)|;$$

(d) $|(T^R)'(x)| > 4^6$;

(e) $m\{x \in \Lambda^{(j)} : R(x) > n\} \leq C''\theta^n$ for some $\theta < 1$ and C'' independent of δ .

Proof. Since $\Lambda^{(j)}$ is an interval of the form Ω in Proposition 4.1, there exists a partition \mathcal{Z}_1 of $\Lambda^{(j)}$ and a stopping time S_1 with the properties of that proposition. For each $\omega_1 \in \mathcal{Z}_1$, we do the following.

Case 1. $\hat{T}^{S_1}\omega_1 \in H$. We set $R(x) = S_1(x)$ for $x \in \omega_1$ and enter ω_1 as an element of the partition η .

Case 2. $|\hat{T}^{S_1}\omega_1| \geq 4^8\varepsilon$. In this case, $\hat{T}^{S_1}\omega_1$ must completely cover at least 4^7 of the $\Lambda^{(i)}$, plus at most one extra piece on each side. If the left-most end piece has length less than ε , then we adjoin it to the left-most $\Lambda^{(i)}$ that is covered by $\hat{T}^{S_1}\omega_1$; otherwise we leave it. We do the same for the right end piece. For each of the $\Lambda^{(i)}$ that has not been adjoined to the end pieces, we enter $\omega_1 \cap T^{-S_1}\Lambda^{(i)}$ as an element of η and declare $R(x) = S_1(x)$ on this interval.

For each $\omega_1 \in \mathcal{Z}_1$ with $|\hat{T}^{S_1}\omega_1| \geq 4^8\varepsilon$, we are left with at most two pieces ω_1^\pm with $\varepsilon \leq |\hat{T}^{S_1}\omega_1^\pm| \leq 3\varepsilon$ on which R has not yet been declared. We apply Proposition 4.1 to obtain a partition and stopping time S on each interval $\hat{T}^{S_1}\omega_1^\pm$. Define $S_2 = S_1 + S \circ \hat{T}^{S_1}$ on ω_1^\pm . This induces a partition \mathcal{Z}_2 on $\Lambda^{(j)} \setminus \{x : R(x) = S_1(x)\}$. For each piece $\omega_2 \in \mathcal{Z}_2$ we apply Cases 1 and 2 described above and as before, are left with two pieces $\hat{T}^{S_2}\omega_2^\pm$ each with length between ε and 3ε such that R has not yet been declared on ω_2^\pm . We use Proposition 4.1 to define $S_3 = S_2 + S \circ \hat{T}^{S_2}$ on $\Lambda^{(j)} \setminus (\{R = S_1\} \cup \{R = S_2\})$ and proceed inductively.

Continuing in this way, we generate a sequence of stopping times S_i and partitions \mathcal{Z}_i of $\Lambda^{(j)} \setminus (\cup_{k=1}^i \{R = S_k\})$ such that S_k is constant on $\omega_i \in \mathcal{Z}_i$ for each $k \leq i$. It is clear that R and η as constructed above satisfy items (a)–(d) by Proposition 4.1. We now derive the tail estimate (e).

Fix an $\omega_i \in \mathcal{Z}_i$ and let φ_i denote the inverse of \hat{T}^{S_i} restricted to ω_i . Then part (e) of Proposition 4.1 yields

$$m\{x \in \hat{T}^{S_i}\omega_i^\pm : S_{i+1}(\varphi_i x) - S_i(\varphi_i x) > n\} \leq C'e^{-n/21}.$$

Using the distortion bound of Lemma 4.6, this becomes

$$m\{x \in \omega_i^\pm : S_{i+1}(x) - S_i(x) > n\} \leq eC' \frac{|\omega_i^\pm|}{|\hat{T}^{S_i}\omega_i^\pm|} e^{-n/21} \leq eC' \frac{|\omega_i^\pm|}{\varepsilon} e^{-n/21}. \quad (15)$$

Since we return at least $1 - 6/4^8$ of $\hat{T}^{S_{i+1}}\omega_i^\pm$, we conclude, again using distortion bounds, that at least $\frac{1}{3}$ of ω_i^\pm is returned at time S_{i+1} , i.e.

$$m\{x \in \omega_i^\pm : R(x) = S_{i+1}(x)\} \geq \frac{|\omega_i^\pm|}{3}. \quad (16)$$

We wish to estimate $m\{x \in \Lambda^{(j)} : R(x) > n\}$. Let $\alpha > 0$ be a small number to be chosen later. So

$$m\{R(x) > n\} = \sum_{i \leq [\alpha n]} m\{S_{i-1} \leq n < S_i\} + \sum_{i > [\alpha n]} m\{S_{i-1} \leq n < S_i\}. \quad (17)$$

Using equation (16) and summing over pieces ω_i^\pm in $\Lambda^{(j)}$, the second sum can be estimated by

$$\sum_{i > [\alpha n]} m\{S_{i-1} \leq n < S_i \leq R(x)\} \leq \sum_{i > [\alpha n]} \left(\frac{2}{3}\right)^{i-1} |\Lambda^{(j)}| = \left(\frac{2}{3}\right)^{[\alpha n]} 3|\Lambda^{(j)}|. \quad (18)$$

To estimate the first sum in equation (17), we define $A_i(l_1, \dots, l_{i-1}) = \{x \in \Lambda^{(j)} : R(x) > n, S_{i-1} \leq n < S_i \text{ and } S_k - S_{k-1} = l_k, k = 1, \dots, i-1\}$. Each term in the first sum can be estimated by

$$m\{R(x) > n : S_{i-1} \leq n < S_i\} = \sum_{\substack{(l_1, \dots, l_{i-1}) \\ \sum_k l_k \leq n}} m(A_i(l_1, \dots, l_{i-1})). \quad (19)$$

For a fixed (l_1, \dots, l_{i-1}) let $B_m = \{x \in \Lambda^{(j)} : R(x) > S_m(x) \text{ and } S_k - S_{k-1} = l_k, k = 1, \dots, m\}$. We condition on the B_m to obtain

$$m(A_i(l_1, \dots, l_{i-1})) = m\{R > n \geq S_{i-1} | B_{i-1}\} \prod_{m=1}^{i-1} m\{S_m - S_{m-1} = l_m | B_{m-1}\}.$$

We estimate this product using equation (15):

$$\begin{aligned} m(A_i(l_1, \dots, l_{i-1})) &\leq eC' \frac{|\Lambda^{(j)}|}{\varepsilon} e^{-(n - \sum_k l_k)/21} \prod_{k=1}^{i-1} eC' \frac{|\Lambda^{(j)}|}{\varepsilon} e^{-(l_k - 1)/21} \\ &\leq (2eC')^i e^{i/21} e^{-n/21}. \end{aligned} \quad (20)$$

Now we estimate

$$\#\left\{(l_1, \dots, l_{i-1}) : \sum_k l_k \leq n\right\} \leq \binom{n}{i-1} \leq (1 + \tau(\alpha))^n,$$

where $\tau(\alpha) \rightarrow 0$ as $\alpha \rightarrow 0$. Using equations (18)–(20), equation (17) becomes

$$\begin{aligned} m\{R(x) > n\} &\leq \sum_{i \leq [\alpha n]} (1 + \tau(\alpha))^n (2eC')^i e^{i/21} e^{-n/21} + \left(\frac{2}{3}\right)^{[\alpha n]} 3|\Lambda^{(j)}| \\ &\leq (1 + \tau(\alpha))^n (2eC')^{\alpha n + 1} e^{(\alpha n + 1)/21} e^{-n/21} + \left(\frac{2}{3}\right)^{[\alpha n]} 3|\Lambda^{(j)}| \\ &\leq C'' \theta^n, \end{aligned}$$

where $\theta := \left(\frac{2}{3}\right)^\alpha$ for the optimal α which makes $\left(\frac{2}{3}\right)^\alpha = (1 + \tau(\alpha))(2eC')^\alpha e^{(\alpha - 1)/21}$. \square

4.4. *Estimating the amount that falls in the hole.* For each $\Lambda^{(j)}$, we estimate the amount of Lebesgue measure that can fall into the hole H at a given time n . This estimate will resemble the estimates of §§4.2 and 4.3. We prove the following proposition.

PROPOSITION 4.3. *There exists a $D > 0$ such that for any $\Lambda^{(j)}$,*

$$m\{x \in \Lambda^{(j)} : R(x) = n \text{ and } \hat{T}^n x \in H\} \leq Dm(H)\theta^n.$$

The proof of this proposition will depend on the following lemma in much the same way that the proof of Proposition 4.2 used Proposition 4.1.

LEMMA 4.4. *Let Ω , \mathcal{Z} and S be as in the statement of Proposition 4.1. There exists a $D' > 0$ such that*

$$m\{x \in \Omega : S(x) = n \text{ and } \hat{T}^n x \in H\} \leq D'm(H)e^{-n/21}.$$

Proof. Suppose $\omega \in \mathcal{Z}$ and $\hat{T}^S \omega \subset H$ with $S = n$. If $\omega \subset I_k$, we set $r_0 = k$ and $p_0 = p(r_0)$; otherwise, set $r_0 = p_0 = 0$. Suppose ω has itinerary (r_1, \dots, r_s) at times t_1, \dots, t_s and let (p_1, \dots, p_s) be the associated recovery times. Note that since $\hat{T}^n \omega \subset H$, ω must be free at time n so that $n \geq t_s + p_s$. Using equation (6), we obtain

$$|\omega| \leq \frac{mH}{c_0 c'_0 \delta} e^{-\frac{1}{3}\lambda_0 n}. \quad (21)$$

Since $\hat{T}^n \omega$ is free, we have $n \geq p_0 + \sum_{i=1}^s p_i \geq \frac{1}{2}(|r_0| + \sum_{i=1}^s |r_i|)$ and $s \leq K/\Delta$, where $K = \sum_{i=1}^s |r_i|$. We estimate using equations (3) and (21):

$$\begin{aligned} m\{x \in I_{r_0} : \hat{T}^n x \in H\} &\leq \sum_{K=0}^{2n} \sum_{s=0}^{K/\Delta} \left(\begin{array}{c} \# \text{ } s\text{-tuples with} \\ \sum |r_i| = K \end{array} \right) \cdot \left(\begin{array}{c} \# \text{ pieces with} \\ \text{same itinerary} \end{array} \right) \cdot \left(\begin{array}{c} \text{measure} \\ 1 \text{ piece} \end{array} \right) \\ &\leq \sum_{K=0}^{2n} \sum_{s=0}^{K/\Delta} 2^s \binom{K-1}{s-1} \cdot 2^{s+1} 2^{(n-(K/2))/m_0} \cdot \frac{mH}{c_0 c'_0 \delta} e^{-\frac{1}{3}\lambda_0 n} \\ &\leq \sum_{K=0}^{2n} 2(1 + \sigma(\delta))^K 4^{(K/\Delta)+1} 2^{-K/2m_0} 2^{n/m_0} \frac{mH}{c_0 c'_0 \delta} e^{-\frac{1}{3}\lambda_0 n} \\ &\leq \frac{mH}{c_0 c'_0 \delta} 8(1 + \sigma(\delta))^{2n+1} 2^{n/m_0} e^{-\frac{1}{4}\lambda_0 n} e^{-\frac{1}{12}\lambda_0 |r_0|} \\ &\leq c_8 m(H) e^{-\frac{1}{21}n} e^{-\frac{1}{12}\lambda_0 |r_0|} \end{aligned}$$

for δ small enough and $m_0 = 10$. But since δ has already been chosen small enough to make a comparable series converge in §4.2, the same δ will work here.

If $\Omega \subset I \setminus (-\delta, \delta)$, then $r_0 = 0$ and the above estimate shows that

$$m\{x \in \Omega : S(x) = n \text{ and } \hat{T}^n x \in H\} \leq c_8 m(H) e^{-\frac{1}{21}n}.$$

If $\Omega \subset (-\delta, \delta)$, then summing across the I_k , we have

$$m\{x \in \Omega : S(x) = n \text{ and } \hat{T}^n x \in H\} \leq c_8 m(H) e^{-\frac{1}{21}n} \sum_{k=k_0}^{\infty} e^{-\frac{1}{12}\lambda_0 k}.$$

This proves the lemma with

$$D' := \max \left\{ c_8, c_8 \frac{e^{-\frac{1}{12}\lambda_0 k_0}}{1 - e^{-\frac{1}{12}\lambda_0}} \right\}.$$

Note that $D' \sim 1/\delta$. □

Proof of Proposition 4.3. Let $F_n = \{x \in \Lambda^{(j)} : R(x) = n \text{ and } \hat{T}^n x \in H\}$. Each $x \in F_n$ will have a number of auxiliary stopping times defined by the construction described in §4.3. We define $S_*(x)$ to be the time when x starts its final auxiliary stopping time before falling into the hole. Let $F_n^i = \{x \in F_n : S_*(x) = i\}$. Note that if $x \in F_n^i$ then $x \in \omega_*^\pm$ and $T^{S_*(x)}\omega_*^\pm$ is an interval of the type Ω in Lemma 4.4. Thus

$$m\{y \in T^{S_*(x)}\omega_*^\pm : S(y) = k \text{ and } \hat{T}^k(y) \in H\} \leq D'm(H)e^{-k/21}.$$

Using the distortion bound of Lemma 4.6, we obtain

$$\begin{aligned} m\{x \in \omega_*^\pm : \hat{T}^n(x) \in H\} &\leq eD'mH \frac{|\omega_*^\pm|}{|T^{S_*(x)}\omega_*^\pm|} e^{-(n-S_*(x))/21} \\ &\leq eD'mH \frac{|\omega_*^\pm|}{\varepsilon} e^{-(n-i)/21}. \end{aligned} \quad (22)$$

Using equation (22) and Proposition 4.2, we estimate the size of F_n :

$$\begin{aligned} m(F_n) &= \sum_{i=0}^{n-1} m(F_n^i) = \sum_{i=0}^{n-1} m\{\hat{T}^n \in H | S_*(x) = i\} \cdot m\{S_*(x) = i\} \\ &\leq \sum_{i=0}^{n-1} 2eD'm(H)e^{-(n-i)/21} \cdot C''\theta^i \leq 2eD'C''m(H)\theta^n \sum_{i=1}^{\infty} \frac{1}{(e^{\frac{1}{21}}\theta)^i} \\ &=: Dm(H)\theta^n. \end{aligned}$$

Note again that $D \sim 1/\delta$. □

4.5. Distortion bounds. We begin this section by deriving the distortion bound on which Properties P1 are based. The content of this estimate is essentially to show that the derivative $(\hat{T}^n)'(\hat{T}x)$ for $x \in (-\delta, \delta)$ is comparable to $(\hat{T}^n)'(\hat{T}0)$ for $0 \leq n \leq p(x) - 1$ and so grows exponentially. We estimate

$$\begin{aligned} \log \frac{(\hat{T}^{n-1})'(\hat{T}x)}{(\hat{T}^{n-1})'(\hat{T}0)} &\leq \sum_{j=1}^{n-1} |\log|\hat{T}^j(x)| - \log|\hat{T}^j(0)|| \\ &\leq \frac{1}{\delta_0} \sum_{j=1}^{n-1} |\hat{T}^j(x) - \hat{T}^j(0)| \\ &\leq \frac{1}{\delta_0} \sum_{j=1}^{n-1} \frac{1}{j^2}. \end{aligned}$$

So we have

$$d_0^{-1}(1.9)^n \leq |(\hat{T}^n)'(\hat{T}x)| \leq d_0(1.9)^n \quad (23)$$

for some $d_0 > 1$ for each $0 \leq n \leq p(x) - 1$. We will use this estimate in §5.4 when determining an upper bound for the conditionally invariant density. The remaining results in this section concern the time $q(k)$ and the main distortion bound for pieces which return at times S and R .

Condition A1 implies that for each $k \geq k_0$ the interval $(0, e^{-k})$ must grow to at least length r before intersecting H . Recall $q(k)$ as defined in §2.4. The following lemma allows us to conclude that each I_k must grow to a fixed length by time $q(k)$.

LEMMA 4.5. *There exists a $c_1 > 0$ independent of δ such that for any k with $k \geq k_0$ and any n with $M_0 < n \leq q(k)$:*

(a) *if $x, y \in (0, e^{-k})$, then*

$$\log \frac{(\hat{T}^{n-1})'(\hat{T}x)}{(\hat{T}^{n-1})'(\hat{T}y)} \leq c_1;$$

(b) *if $x, y \in I_k$, then*

$$\log \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} \leq c_1.$$

Proof. Write $n - 1 = lM_0 + j$ for some $l \in \mathbb{Z}^+$, $0 \leq j \leq M_0 - 1$. For $x, y \in I_k$, we estimate the derivatives using Definition 2.1(b)(i):

$$\begin{aligned} \log \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} &\leq \left| \log \frac{x}{y} \right| + \sum_{i=0}^{l-2} |\log|(\hat{T}^{M_0})'(\hat{T}^{1+iM_0}x)| - \log|(\hat{T}^{M_0})'(\hat{T}^{1+iM_0}y)|| \\ &\quad + |\log|(\hat{T}^{M_0+j})'(\hat{T}^{n-(M_0+j)}x)| - \log|(\hat{T}^{M_0+j})'(\hat{T}^{n-(M_0+j)}y)|| \\ &\leq \left| \log \frac{x}{y} \right| + \sum_{i=0}^{l-2} \frac{1}{e^{\lambda_0 M_0}} |(\hat{T}^{M_0})'(\hat{T}^{1+iM_0}x) - (\hat{T}^{M_0})'(\hat{T}^{1+iM_0}y)| \\ &\quad + \frac{1}{e^{\lambda_0(M_0+j)}} |(\hat{T}^{M_0+j})'(\hat{T}^{n-M_0-j}x) - (\hat{T}^{M_0+j})'(\hat{T}^{n-M_0-j}y)| \\ &\leq \left| \log \frac{x}{y} \right| + \sum_{i=0}^{l-2} \frac{(2a)^{M_0}}{e^{\lambda_0 M_0}} |\hat{T}^{1+iM_0}(x) - \hat{T}^{1+iM_0}(y)| \\ &\quad + \frac{(2a)^{M_0+j}}{e^{\lambda_0(M_0+j)}} |\hat{T}^{n-M_0-j}(x) - \hat{T}^{n-M_0-j}(y)| \\ &\leq \left| \log \frac{x}{y} \right| + \frac{(2a)^{M_0}}{e^{\lambda_0 M_0}} \sum_{i=0}^{l-2} |\hat{T}^n(x) - \hat{T}^n(y)| e^{-\lambda_0((l-i)M_0+j)} \\ &\quad + \frac{(2a)^{M_0+j}}{e^{\lambda_0(M_0+j)}} |\hat{T}^n(x) - \hat{T}^n(y)| e^{-\lambda_0(M_0+j)} \\ &\leq \left| \log \frac{x}{y} \right| + |\hat{T}^n(x) - \hat{T}^n(y)| \frac{(2a)^{M_0}}{e^{\lambda_0 M_0}} \sum_{i=2}^{\infty} e^{-\lambda_0 M_0 i} \\ &\quad + |\hat{T}^n(x) - \hat{T}^n(y)| \frac{(2a)^{2M_0-1}}{e^{2\lambda_0(2M_0-1)}} \end{aligned} \tag{24}$$

$$\leq 1 + \frac{\delta_0}{2} \left(\frac{(2a)^{M_0}}{e^{\lambda_0 M_0}} \sum_{i=2}^{\infty} e^{-\lambda_0 M_0 i} + \frac{(2a)^{2M_0-1}}{e^{2\lambda_0(2M_0-1)}} \right)$$

$$=: c_1.$$

This proves (b) directly and (a) follows simply by omitting the first term of the sum. \square

Lemma 4.5(a) says that for any $k \geq k_0$, the relative scale of the partition $\{\hat{T}(I_j)\}_{j \geq k}$ of $\hat{T}((0, e^{-k}))$ is maintained until time $q(k)$. The same is also true for $\hat{T}((-e^{-k}, 0))$. Since the ratio of $|\hat{T}((0, e^{-k}))|$ to $|\hat{T}((0, e^{-k-1}))|$ is e^2 , the ratio of $|\hat{T}^q((0, e^{-k}))|$ to $|\hat{T}^q((0, e^{-k-1}))|$ is about e^2 as well. Also $|\hat{T}^q((0, e^{-k}))| \geq r/8$ implies that $|\hat{T}^q(I_k)|$ is uniformly bounded below. This minimum length is the quantity ε' introduced in §2.4.

Lemma 4.5(b) yields a distortion bound for $x, y \in I_k$ at time $q(k)$:

$$\frac{|\hat{T}^q(x) - \hat{T}^q(y)|}{|x - y|} \geq e^{-c_1} \frac{|\hat{T}^q I_k|}{|I_k|} \geq e^{-c_1} \frac{\varepsilon'}{|I_k|}.$$

This implies that

$$|x - y| \leq e^{c_1} \frac{|I_k|}{\varepsilon'} |\hat{T}^q(x) - \hat{T}^q(y)|.$$

Now we substitute the above estimate into equation (24) to obtain

$$\begin{aligned} \log \frac{(\hat{T}^q)'(x)}{(\hat{T}^q)'(y)} &\leq \frac{|x - y|}{e^{-k-1}} + |\hat{T}^q x - \hat{T}^q y| \left(\frac{(2a)^{M_0}}{e^{\lambda_0 M_0}} \sum_{i=2}^{\infty} e^{-\lambda_0 M_0 i} + \frac{(2a)^{2M_0-1}}{e^{2\lambda_0(2M_0-1)}} \right) \\ &\leq |\hat{T}^q x - \hat{T}^q y| \left(e^{c_1} \frac{2}{\varepsilon'} + \frac{(2a)^{M_0}}{e^{\lambda_0 M_0}} \sum_{i=2}^{\infty} e^{-\lambda_0 M_0 i} + \frac{(2a)^{2M_0-1}}{e^{2\lambda_0(2M_0-1)}} \right) \\ &=: c_2 |\hat{T}^q x - \hat{T}^q y|. \end{aligned}$$

From this we conclude that for $x, y \in I_k$,

$$\left| \frac{(\hat{T}^q)'(x)}{(\hat{T}^q)'(y)} - 1 \right| \leq c_2 |\hat{T}^q(x) - \hat{T}^q(y)|. \quad (25)$$

Equation (25) allows us to prove our main distortion lemma for an interval ω which is returned at time n .

LEMMA 4.6. (Distortion bounds) *If an interval $\omega \subset I$ is such that $\hat{T}\omega$ lies in one element of \mathcal{Q} for each $i \in [0, n]$ and $S(\omega) = n$, then there exists a constant $\tilde{C} > 0$ such that*

$$\left| \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} - 1 \right| \leq \tilde{C} |\hat{T}^n(x) - \hat{T}^n(y)|.$$

Proof. Let $t_0 = 0$, $t_{s+1} = n$ and t_1, \dots, t_s be the times that ω visits $(-\delta, \delta)$ before time n . $\hat{T}^{t_i} \omega \subset I_{k_i}$ so we set $q_i = q(k_i)$. Since $S(\omega) = n$, we know that $n > t_s + q_s$ so for each i

we can write $t_{i+1} = t_i + q_i + l_i$ for some $l_i \in \mathbb{N}$. Then using equation (25), we estimate

$$\begin{aligned} \log \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} &= \log \prod_{i=0}^s \frac{(\hat{T}^{q_i})'(\hat{T}^{t_i}x) \cdot (\hat{T}^{l_i})'(\hat{T}^{t_i+q_i}x)}{(\hat{T}^{q_i})'(\hat{T}^{t_i}y) \cdot (\hat{T}^{l_i})'(\hat{T}^{t_i+q_i}y)} \\ &= \sum_{i=0}^s \log \frac{(\hat{T}^{q_i})'(\hat{T}^{t_i}x)}{(\hat{T}^{q_i})'(\hat{T}^{t_i}y)} + \log \frac{(\hat{T}^{l_i})'(\hat{T}^{t_i+q_i}x)}{(\hat{T}^{l_i})'(\hat{T}^{t_i+q_i}y)} \\ &\leq \sum_{i=0}^s c_2 |\hat{T}^{t_i+q_i}x - \hat{T}^{t_i+q_i}y| + \log \prod_{j=0}^{l_i-1} \frac{\hat{T}'(\hat{T}^{t_i+q_i+j}x)}{\hat{T}'(\hat{T}^{t_i+q_i+j}y)}. \end{aligned} \quad (26)$$

We estimate the second term by

$$\begin{aligned} \log \prod_{j=0}^{l_i-1} \frac{\hat{T}'(\hat{T}^{t_i+q_i+j}x)}{\hat{T}'(\hat{T}^{t_i+q_i+j}y)} &\leq \sum_{j=0}^{l_i-1} |\log \hat{T}'(\hat{T}^{t_i+q_i+j}x)| - |\log \hat{T}'(\hat{T}^{t_i+q_i+j}y)| \\ &\leq \sum_{j=0}^{l_i-1} \frac{1}{2a\delta} |\hat{T}'(\hat{T}^{t_i+q_i+j}x) - \hat{T}'(\hat{T}^{t_i+q_i+j}y)| \\ &\leq \sum_{j=0}^{l_i-1} \frac{1}{\delta} |\hat{T}^{t_i+q_i+j}(x) - \hat{T}^{t_i+q_i+j}(y)|. \end{aligned}$$

We substitute this result back into equation (26) and use Lemma 2.5(a) to get

$$\begin{aligned} \log \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} &\leq \sum_{i=0}^s \left(\frac{c_2}{c'_0\delta} e^{-\frac{1}{3}\lambda_0 l_i} + \sum_{j=0}^{l_i-1} \frac{1}{\delta} \frac{1}{c'_0\delta} e^{-\frac{1}{3}\lambda_0(l_i-j)} \right) |\hat{T}^{t_{i+1}}(x) - \hat{T}^{t_{i+1}}(y)| \\ &\leq \sum_{i=0}^s \left(c_2 \frac{1}{c'_0\delta} + \frac{1}{\delta} \frac{1}{c'_0\delta} \sum_{j=1}^{\infty} e^{-\frac{1}{3}\lambda_0 j} \right) |\hat{T}^{t_{i+1}}(x) - \hat{T}^{t_{i+1}}(y)| \\ &=: c_3 \sum_{i=0}^s |\hat{T}^{t_{i+1}}(x) - \hat{T}^{t_{i+1}}(y)|. \end{aligned} \quad (27)$$

We estimate $|\hat{T}^{t_i}(x) - \hat{T}^{t_i}(y)|$ using equation (5). Since $|(\hat{T}^{t_{i+1}-t_i})'(\hat{T}^{t_i}x)| \geq e^{\frac{1}{3}\lambda_0(t_{i+1}-t_i)}$, we have

$$|\hat{T}^{t_i}(x) - \hat{T}^{t_i}(y)| \leq e^{-\frac{1}{3}\lambda_0(t_{i+1}-t_i)} |\hat{T}^{t_{i+1}}(x) - \hat{T}^{t_{i+1}}(y)| \leq e^{-\frac{1}{3}\lambda_0(n-t_i)} |\hat{T}^n(x) - \hat{T}^n(y)|.$$

We substitute this back into equation (27) to conclude the proof:

$$\begin{aligned} \log \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} &\leq c_3 \sum_{i=0}^s e^{-\frac{1}{3}\lambda_0(n-t_{i+1})} |\hat{T}^n(x) - \hat{T}^n(y)| \\ &\leq c_3 |\hat{T}^n(x) - \hat{T}^n(y)| \sum_{i=0}^{\infty} e^{-\frac{1}{6}\lambda_0 k_0 i} \\ &=: \tilde{C} |\hat{T}^n(x) - \hat{T}^n(y)|. \end{aligned} \quad \square$$

Note that the constant $\tilde{C} \sim 1/\delta^2$. Also, this Lemma proves item (c) of Proposition 4.1.

Item (d) of Proposition 4.1 follows immediately using the assumption that $\varepsilon \leq 1/4^9 \tilde{C}$ and noting that $|\hat{T}^S x - \hat{T}^S y| < 4^9 \varepsilon$ since $|T^{S-1} \omega| \leq 4^8 \varepsilon$ and $|\hat{T}'| \leq 4$. Since $|T^S \omega|/|\omega| \geq 4^8/3$, for any $x \in \omega$ we must have

$$|(T^S)'(x)| \geq \frac{4^8}{3} e^{-\tilde{C}4^9 \varepsilon}$$

from which $|(T^S)'(x)| > 4^6$ follows.

Remark. The weaker bound

$$\left| \frac{(\hat{T}^n)'(x)}{(\hat{T}^n)'(y)} - 1 \right| \leq \tilde{C} \quad (28)$$

can be proved using Lemma 4.5(b) instead of equation (25) in equation (26). Although the bound is weaker than that in Lemma 4.6, it is valid for any time n when $\hat{T}^n \omega$ is free, not just when $n = S(\omega)$. We shall use this bound later in §5.4.

5. An a.c.c.i.m. for the logistic map

5.1. Defining the tower map. We identify $\Lambda^{(1)}, \dots, \Lambda^{(N)}$ with N intervals of unit length, $\hat{\Delta}_0^{(1)}, \dots, \hat{\Delta}_0^{(N)}$. The partition η and return time function R of Proposition 4.2 induce a partition and return time function on $\hat{\Delta}_0 := \bigcup_i \hat{\Delta}_0^{(i)}$ which we refer to by the same names. We define the tower as usual:

$$\hat{\Delta} := \{(x, n) \in \hat{\Delta}_0 \times \mathbb{N} : n < R(x)\}.$$

Recall the notation $\hat{\Delta}_l = \hat{\Delta}_{|n=l}$ for the l th level of the tower, and let $\hat{\Delta}_l^{(i)}$ denote the l th level above $\hat{\Delta}_0^{(i)}$. Define the projection $\pi_0 : \hat{\Delta}_0 \rightarrow I$ as piecewise linear with $\pi_0(\hat{\Delta}_0^{(i)}) = \Lambda^{(i)}$ and $\pi'_0|_{\hat{\Delta}_0^{(i)}} = |\Lambda^{(i)}|$. This makes $\varepsilon \leq \pi'_0 \leq 2\varepsilon$ on $\hat{\Delta}_0$. If $\hat{T}^R \omega \subset H$, then we put a hole $\tilde{H}_{R(\omega),j}^{(i)}$ in the $R(\omega)$ level of the tower above $\pi_0^{-1} \omega$ in $\hat{\Delta}_0^{(i)}$. This defines the hole \tilde{H} in the tower $\hat{\Delta}$. Let $\Delta = \hat{\Delta} \setminus \tilde{H}$ and in general $\Delta_l^{(i)} = \hat{\Delta}_l^{(i)} \setminus \tilde{H}$.

Let \hat{F} be the tower map, $\hat{F} : \Delta \rightarrow \hat{\Delta}$. Define a projection $\pi : \hat{\Delta} \rightarrow [-1, 1]$ such that $\pi \circ \hat{F} = \hat{T} \circ \pi$ on Δ . The elements of the Markov partition $\Delta_{l,j}^{(i)}$ are the maximal intervals on $\hat{\Delta}_l^{(i)}$ which project onto the dynamically defined elements of Ω_l in the construction of the return time function R above the reference interval $\Lambda^{(i)}$, with the exception that each $\Delta_0^{(i)}$ is taken to be a single element of the partition. Recall that $\Delta_{l,j}^{(i)*}$ are the elements which return to Δ_0 at time $l+1$.

For $x \in \Delta$, we have the identity $\pi' \circ \hat{F}^l(x) \cdot (\hat{F}^l)'(x) = (\hat{T}^l)' \circ \pi(x) \cdot \pi'(x)$. If $R(x) > l$ then $(\hat{F}^l)'(x) = 1$ so

$$\pi' \circ \hat{F}^l(x) = (\hat{T}^l)' \circ \pi(x) \cdot \pi'(x). \quad (29)$$

Now let $z \in \Delta_0^{(i)}$ with $R(z) = l+1$. Let $\hat{F}^l z = \bar{x}$. Then $\pi' \circ \hat{F}^{l+1}(z) \cdot (\hat{F}^{l+1})'(z) = (\hat{T}^{l+1})' \circ \pi(z) \cdot \pi'(z)$. But $\hat{F}^{l+1}(z) \in \Delta_0^{(j)}$ so that $\pi' \circ \hat{F}^{l+1}(z) = |\Lambda^{(j)}|$ and $(\hat{F}^{l+1})'(z) = \hat{F}'(\hat{F}^l z) \cdot (\hat{F}^l)'(z) = \hat{F}'(\bar{x})$. This yields

$$\hat{F}'(\bar{x}) = (T^{l+1})'(\pi z) \frac{|\Lambda^{(i)}|}{|\Lambda^{(j)}|}. \quad (30)$$

Using this fact, Proposition 4.2(d) and equation (6), we conclude that

$$\inf_{\Delta^*} |\hat{F}'| > \frac{4^6}{2} \quad \text{and} \quad \inf_{\Delta_l^*} |\hat{F}'| > \frac{c_0 c_0' \delta}{2} e^{\frac{1}{3} \lambda_0 (l+1)}. \quad (31)$$

We derive a distortion estimate for \hat{F} . Let $x, y \in \Delta_{l,j}^{(k)}$ be such that $\hat{F}x, \hat{F}y \in \hat{\Delta}_0^{(i)}$. Using Lemma 4.6 and equation (30) we have

$$\begin{aligned} \left| \frac{\hat{F}'(x)}{\hat{F}'(y)} - 1 \right| &= \left| \frac{(\hat{T}^{l+1})'(\pi \circ \hat{F}^{-l}x)}{(\hat{T}^{l+1})'(\pi \circ \hat{F}^{-l}y)} - 1 \right| \\ &\leq \tilde{C} |\hat{T}^{l+1}(\pi \circ \hat{F}^{-l}x) - \hat{T}^{l+1}(\pi \circ \hat{F}^{-l}y)| \\ &= \tilde{C} |\pi \circ \hat{F}(x) - \pi \circ \hat{F}(y)| \\ &\leq \tilde{C}(2\varepsilon) |\hat{F}(x) - \hat{F}(y)|. \end{aligned}$$

Let $F = \hat{F}|(\Delta \setminus \hat{F}^{-1}\tilde{H})$. F also satisfies the relation $\pi \circ F = T \circ \pi$ on its domain, and so the above estimates hold for F' .

5.2. (Δ, F, m) satisfies Conditions H1 and H2. Recall Properties P2(a)–(c) required of the tower map in §3 as well as Conditions H1 and H2. It is clear from the discussion of the previous section that \hat{F} has properties (a) and (c) with $C = 2\tilde{C}\varepsilon \leq 2/4^9$. Property (b) follows from equation (31). Let ν be the smallest integer such that $(c_0 c_0' \delta / 2) e^{\frac{1}{6} \lambda_0 (\nu+1)} \geq 2$. For $l \geq \nu$, $|F'| \geq 2e^{\frac{1}{6} \lambda_0 (l+1)}$, while for $l < \nu$, $|F'| > 2(4^5)^{l/\nu}$. So (b) follows with $\gamma = 2$ and $\beta = \min\{\lambda_0/6, (5/\nu) \log 4\}$.

Condition H1 is satisfied with the same θ as in the statement of Proposition 4.2. This is because $m\hat{\Delta}_n = m\Delta_n + m\tilde{H}_n$ and Proposition 4.2 yields $m\Delta_n \leq (NC''/\varepsilon)\theta^n$ while Proposition 4.3 yields $m\tilde{H}_n \leq (NDmH/\varepsilon)\theta^n$.

In §3, ξ is defined so that $e^{-\xi} > \max\{\theta, e^{-\beta}\}$. Actually, from the proof of Proposition 4.2, the rate of contraction of θ is slower than $e^{-\beta}$ so we may choose $\xi = -\frac{1}{2} \log \theta$. Then Condition H2 becomes

$$\sum_{l=1}^{\infty} \theta^{-(l-1)/2} m\tilde{H}_l < \frac{(1 - \sqrt{\theta})^2}{1 + C}.$$

But $m\tilde{H}_l \leq (NDmH/\varepsilon)\theta^l$. So

$$\sum_{l=1}^{\infty} \theta^{-(l-1)/2} m\tilde{H}_l \leq \sum_{l=1}^{\infty} \frac{NDmH}{\varepsilon} \theta^{-(l-1)/2} \theta^l \leq \frac{NDmH}{\varepsilon} \frac{\theta}{1 - \sqrt{\theta}}.$$

Using this estimate, we see that Condition H2 will be satisfied if H satisfies

$$mH < \frac{(1 - \sqrt{\theta})^2}{1 + C} \cdot \frac{\varepsilon(1 - \sqrt{\theta})}{ND\theta},$$

which is slightly weaker than Condition A3.

5.3. *Existence and lower bound for an a.c.c.i.m.* Since (Δ, F, m) satisfies Condition H1 and Condition H2, we conclude that there exists $\varphi \in X_M$ such that $d\tilde{\mu} := \varphi dm$ is an a.c.c.i.m. with respect to F acting on Δ . Let λ be the eigenvalue of $\tilde{\mu}$ and note that $\lambda \geq \sqrt{\theta}$ by the remark following Theorem 3.1. By that same remark we have

$$\lambda \geq 1 - M \sum_{l \geq 1} e^{\xi(l-1)} m(\tilde{H}_l) \geq 1 - \frac{ND\theta}{\varepsilon(1 - \sqrt{\theta})} m(H) \quad (32)$$

by the estimates of §5.2 based on Condition A3.

Now define a measure μ on I by

$$\mu(A) := \tilde{\mu}(\pi^{-1}A)$$

for any Borel subset A of I . Then μ will be an a.c.c.i.m. with respect to T with the same eigenvalue λ since for any Borel $A \subset I$,

$$\begin{aligned} \mu(T^{-1}A) &:= \tilde{\mu}(\pi^{-1} \circ T^{-1}A) = \tilde{\mu}(F^{-1} \circ \pi^{-1}A) \\ &= \lambda \tilde{\mu}(\pi^{-1}A) =: \lambda \mu(A). \end{aligned}$$

Let ψ be the density of the measure μ . The fact that ψ is bounded away from zero relies on the genericity Condition A4 as well as the following lemma which is proved in a more general case in [Y1] as Lemma 2.1.

LEMMA 5.1. *Let $\hat{T} \in \mathcal{M}$. For every interval $J \subseteq [-1, 1]$, there exists $n = n(J)$ such that $\hat{T}^n J \supseteq [\hat{T}^2 0, \hat{T} 0] = [1 - a, 1]$.*

The integer n in the above lemma can be chosen to depend only on the length of the interval J . If we consider only those intervals with length at least $\varepsilon_0/2$, then we can choose a single $n_0 = n_0(\varepsilon_0)$ such that any such interval J satisfies $\hat{T}^{n_0} J \supseteq [1 - a, 1]$. This is the n_0 introduced in §2.2.

For convenience, we recall Condition A4 of §2.2.

Condition A4.

- (a) $\hat{T}^i H_j \cap G_k = \emptyset$ for all $j, k \in [1, \dots, L]$, $0 \leq i \leq n_0$.
- (b) $\hat{T}^i H_j \cap H_k = \emptyset$ for all $j, k \in [1, \dots, L]$, $1 \leq i \leq n_0$.

Recall that the intervals G_j in the statement of Condition A4 are the symmetric counterparts of the H_j so that $\hat{T}^{-1}(\hat{T}H_j) = H_j \cup G_j$.

LEMMA 5.2. *Given any $J \subseteq I$ such that $|J| \geq \varepsilon_0/2$, then*

$$\bigcup_{i=0}^{2n_0} T^i J \supseteq [1 - a, 1] \setminus H.$$

Proof. Fix J as in the statement of the lemma. Suppose there exists an interval ω such that $\omega \cap (\bigcup_{i=0}^{n_0} T^i J) = \emptyset$. Since $\omega \subseteq \hat{T}^{n_0} J$, we must have $\omega \cap \hat{T}^{i_k} H_k \neq \emptyset$ for some H_k such that $H_k \cap \hat{T}^{i'_k} J \neq \emptyset$, for some integers i_k, i'_k with $i_k + i'_k \leq n_0$. In other words, the piece of J that should have covered part of ω fell into H_k before time n_0 . In particular, Condition A4(a) implies that G_1, \dots, G_L are covered by time n_0 , i.e. $G_k \subset T^{n_0} J$.

Condition A4(b) says that G_k cannot fall into the hole again before time n_0 so that $T^{i_k} G_k = \hat{T}^{i_k} H_k$. We conclude that the part of ω which should have been covered by the piece of J that fell into H_k is at the latest covered by an iterate of G_k at time $n_0 + i_k$. Doing this for each k , we have $\omega \subset \bigcup_{k=1}^L T^{i_k} G_k$ and so $\omega \subset \bigcup_{i=0}^{2n_0} T^i J$. \square

In §4.1, we showed that every interval of length at least ε grows to length $4^8 \varepsilon$ in exponential time that depends only on ε . In fact, the construction holds as long as the interval remains less than length ε_0 due to Condition A2. This allows us to conclude that every interval of length ε grows to length ε_0 in exponential time and from there, by Lemma 5.2, it covers $[1 - a, 1]$ by time $2n_0$. This implies that the density ψ is bounded away from zero on $[1 - a, 1] \setminus H$.

5.4. Shape of the density and proof of Theorem 2.4. In this section, we derive bounds on the density ψ and show that it has the form given in the statement of Theorem 2.3. Since the bounds depend only on $H = H_t$ in a sequence of holes of the form described in Theorem 2.4, they are uniform in s and allow us to prove Theorem 2.4.

Let \hat{T} and $\{H_s\}$ be as in the statement of Theorem 2.4. Let $I_s = \hat{I} \setminus H_s$ and $T_s = \hat{T}|_{I_s} \cap \hat{T}^{-1} I_s$. The assumptions on the holes imply that each H_s satisfies Conditions A1–A4 with the same choice of constants. This is because the intervals of monotonicity of the map T_s only increase in length as $s \rightarrow 0$. So we may apply Theorems 2.2 and 2.3 for each s .

Let φ_s be the conditionally invariant density for $\tilde{\mu}_s$ on Δ with eigenvalue λ_s . Let ψ_s be the density for μ_s , the conditionally invariant measure for T_s on I_s . We fix s and show that ψ_s has lower and upper bounds that are independent of s .

Lower Bound. Recall that $\varphi_s \in X_M$ where $M = (1 + C)/(1 - \sqrt{\theta})$. Thus

$$\begin{aligned} 1 &= \sum_{l=0}^{\infty} \int_{\Delta_l} \varphi_s dm \leq \sum_{l=0}^{\infty} \sup_{\Delta_l} \varphi_s m \Delta_l \leq \sup_{\Delta_0} \varphi_s \sum_{l=0}^{\infty} \frac{1}{\lambda_s^l} m \Delta_l \\ &\leq \sup_{\Delta_0} \varphi_s \sum_{l=0}^{\infty} \frac{C'' N}{\varepsilon \lambda_s^l} \theta^l \leq \frac{C''}{\varepsilon^2} \frac{1}{1 - \sqrt{\theta}} \sup_{\Delta_0} \varphi_s \end{aligned}$$

since $\lambda_s \geq \sqrt{\theta}$. This implies that there exists an i such that

$$\sup_{\Delta_0^{(i)}} \varphi_s \geq \frac{\varepsilon^2(1 - \sqrt{\theta})}{C''}.$$

The regularity of φ_s yields a lower bound on the density,

$$\inf_{\Delta_0^{(i)}} \varphi_s \geq \frac{\varepsilon^2(1 - \sqrt{\theta})}{C''(1 + M)},$$

which in turn implies

$$\inf_{\Delta^{(i)}} \psi_s \geq \frac{\varepsilon(1 - \sqrt{\theta})}{2C''(1 + M)}. \quad (33)$$

Since the length scale ε_0 can be chosen independent of s (if ε_0 works for H_t , it will automatically work for each H_s in the sequence), the constant n_0 of Lemma 5.2 is independent of s . The length scale ε is also independent of s so we conclude that $\Lambda^{(i)}$ will grow to cover $[1 - a, 1] \setminus H_s$ in a fixed number of iterates depending only on ε and ε_0 . Call this number N_0 .

For any $x \in [1 - a, 1] \setminus H_s$, there exists $z \in \Lambda^{(i)}$ and $n \leq N_0$ such that $T_s^n(z) = x$. Let \mathcal{P}_s be the Perron–Frobenius operator associated with the map T_s . Then

$$\lambda_s^n \psi_s(x) = \mathcal{P}_s^n \psi_s(x) = \sum_{y \in T_s^{-n}x} \frac{\psi_s(y)}{|(T_s^n)'(y)|} \geq \frac{\psi_s(z)}{|(T_s^n)'(z)|}.$$

So

$$\inf_{[1-a, 1] \setminus H_s} \psi_s \geq \frac{\varepsilon(1 - \sqrt{\theta})}{4^{N_0} 2(1 + M)C''},$$

which is a lower bound independent of s .

Upper Bound. For the upper bound, we first estimate the number of preimages under the projection π a point in I can have on any given level of the tower Δ . To do this, we consider how many unreturned pieces can be generated while iterating one of the reference intervals $\Lambda^{(i)}$. Once a piece is returned, it no longer generates preimages on subsequent levels of the tower. There are several ways that pieces can be generated.

- (1) An interval intersects the hole and is cut into two pieces. This can happen at most once every m_0 iterates by Condition A2.
- (2) An interval grows to length $4^8 \varepsilon$ and the stopping time S is declared. Most of this interval is returned, except for the two end pieces which continue to be iterated. Thus up to two new pieces are formed. Since each piece begins with length less than 3ε and must grow to length $4^8 \varepsilon$ before another stopping time is declared, this can only happen once every eight iterates.
- (3) An interval lands on 0, the critical point. Then we consider that two new pieces are formed, one on each side of 0. This can happen at most once every $p(k_0)$ iterates. Note that $p(k_0) \geq k_0/2 \geq 25$.
- (4) An interval which lands in $(-\delta, \delta)$ reaches its recovery time. Suppose a piece ω is mapped onto an interval extending from I_r to I_s at time t . We label the subinterval of ω which lands in I_k at time t as ω_k . Without loss of generality, assume $0 < s \leq r \leq \infty$. We consider ω as one piece from time t until time $t + q(s)$. At time $t + q(s)$, ω_s is counted as a separate piece. If $|T^{t+q(s)}\omega_s| < 4^8 \varepsilon$, then we simply continue to iterate it. It will generate new pieces at the rate described by items (1)–(3) above until the next time it enters $(-\delta, \delta)$. If $|T^{t+q(s)}\omega_s| \geq 4^8 \varepsilon$, then by definition of the stopping time S after Proposition 4.1 and the stopping time R described after Proposition 4.2, only one new piece will not be returned at time $t + q(s)$ (as opposed to the usual two pieces which are not returned when R is declared on the middle part of an interval). This is because in the construction, the piece of ω_s which does not completely cover the last $\Lambda^{(j)}$ on the side near ω_{s+1} is adjoined to ω_{s+1} and the stopping time S is not declared on this piece until time $t + q(s + 1)$. Thus returns of this type generate at most *linear growth* in the number of pieces which can overlap at any given time.

We see from these considerations that the number of pieces which can overlap at time n and are generated by the single interval $\Lambda^{(i)}$ satisfies the following bound:

$$\#\{\text{Pieces}\} \leq n2^{(n/m_0+n/8+n/p(k_0))+3} \leq 8n2^{53n/200}, \quad (34)$$

where we have used the fact that $m_0 \geq 10$ and $p(k_0) \geq 25$.

We denote by $\pi_{l,j}^{(i)-1}$ the inverse of $\pi|_{\Delta_{l,j}^{(i)}}$. Let $\psi = \psi_s$ and $\varphi = \varphi_s$ for any $s \in [0, t]$. The density ψ can be written as

$$\psi(x) = \sum_{i=1}^N \sum_l \sum_j \frac{\varphi(\pi_{l,j}^{(i)-1}x)}{\pi'(\pi_{l,j}^{(i)-1}x)}.$$

We seek to estimate this sum by determining the growth of $\pi'(\pi_{l,j}^{(i)-1}x)$. In general we have the relation $\pi'(F^l y) \cdot (F^l)'(y) = (T^l)'(\pi y) \cdot \pi'(y)$. Letting $y = F^{-l}(\pi_{l,j}^{(i)-1}x) \in \Delta_0$, we have $(F^l)'(y) = 1$ and $\pi'(y) = |\Lambda^{(i)}|$ so that

$$\pi'(\pi_{l,j}^{(i)-1}x) = (T^l)'(\pi y) \cdot |\Lambda^{(i)}|. \quad (35)$$

Let l_j be the smallest positive integer $k \leq l$ such that $\pi \circ F^{-k}(\pi_{l,j}^{(i)-1}x) \in (-\delta, \delta)$. If no such k exists, then set $l_j = l$ and do Case 1 below.

Case 1. If $l_j \geq p(\pi \circ F^{-l_j}(\pi_{l,j}^{(i)-1}x))$, then $\pi(\Delta_{l,j}^{(i)})$ is free so that $|(T^l)'(\pi y)| \geq c_0 c'_0 \delta e^{\frac{1}{3}\lambda_0 l}$, using equation (6).

Case 2. If $l_j < p(\pi \circ F^{-l_j}(\pi_{l,j}^{(i)-1}x))$, then we estimate $(T^l)'(\pi y)$ as follows:

$$(T^l)'(\pi y) = (\hat{T}^{l_j})'(T^{l-l_j}(\pi y)) \cdot (\hat{T}^{l-l_j})'(\pi y).$$

The second factor in the above expression is greater than or equal to $c_0 c'_0 \delta e^{\frac{1}{3}\lambda_0(l-l_j)}$ since $\hat{T}^{l-l_j-1}(\pi y)$ is free. To estimate the first factor, we use equation (23) of §4.5 to note that for $z \in (-\delta, \delta)$,

$$\begin{aligned} |(\hat{T}^n)'(z)| &= \hat{T}'(z) \cdot (\hat{T}^{n-1})'(\hat{T}z) \geq 2a|z| \cdot \frac{1}{d_0}(1.9)^{n-1} \\ &\geq 2a \sqrt{\frac{|\hat{T}^n z - \hat{T}^n 0|}{d_0(1.9)^{n-1}}} \cdot \frac{1}{d_0}(1.9)^{n-1} = \frac{2a}{d_0^{3/2}}(1.9)^{(n-1)/2} \sqrt{|\hat{T}^n z - \hat{T}^n 0|}. \end{aligned}$$

Thus

$$(T^l)'(\pi y) \geq d'_0(1.9)^{(l_j-1)/2} e^{\frac{1}{3}\lambda_0(l-l_j)} \sqrt{|x - \hat{T}^{l_j} 0|},$$

where $d'_0 = (2a/d_0^{3/2})c_0 c'_0 \delta$.

Let $p_{l,j}^{(i)} = p(\pi \circ F^{-l_j}(\pi_{l,j}^{(i)-1}x))$. Recall that $\varphi \leq M/\lambda_s^l \leq M e^{l/42}$ on Δ_l . Putting Cases 1 and 2 together, we have the following bound on $\psi(x)$:

$$\psi(x) \leq \sum_i \sum_{\Delta_{l,j}^{(i)}: l_j \geq p_{l,j}^{(i)}} \frac{M e^{l/42}}{c_0 c'_0 \delta \varepsilon} e^{-\frac{1}{3}\lambda_0 l} + \sum_i \sum_{\Delta_{l,j}^{(i)}: l_j < p_{l,j}^{(i)}} \frac{2M e^{\frac{l}{42}} e^{-\frac{1}{3}\lambda_0(l-l_j)} (1.9)^{-l_j/2}}{d'_0 \varepsilon \sqrt{|x - \hat{T}^{l_j} 0|}}. \quad (36)$$

Using equation (34), we see that the first sum is less than

$$\sum_i \sum_l \frac{M e^{l/42}}{c_0 c'_0 \delta \varepsilon} e^{-\frac{1}{3} \lambda_0 l} 8 l 2^{53l/200} < \infty$$

if we take $\lambda_0 = \log 1.9$.

We fix i and estimate the second term by

$$\begin{aligned} & \sum_{\substack{\Delta_{l,j}^{(i)} \\ l_j < p_{l,j}^{(i)}}} \frac{2M e^{l/42} e^{-\frac{1}{3} \lambda_0 (l-l_j)} (1.9)^{-l_j/2}}{d'_0 \varepsilon \sqrt{|x - \hat{T}^{l_j} 0|}} \\ & \leq \sum_k \sum_{\Delta_{l,j}^{(i)}: l_j = k} \frac{2M e^{l/42} e^{-\frac{1}{3} \lambda_0 (l-k)} (1.9)^{-k/2}}{d'_0 \varepsilon \sqrt{|x - \hat{T}^k 0|}} \\ & \leq \sum_{k,l} \frac{2M e^{l/42} e^{-\frac{1}{3} \lambda_0 (l-k)} (1.9)^{-k/2}}{d'_0 \varepsilon \sqrt{|x - \hat{T}^k 0|}} 8(l-k) 2^{53(l-k)/200}, \end{aligned}$$

where in the second line we have used the fact that the number of pieces we are summing over from time k to time l has not changed since these pieces are bound during that time.

Using $\lambda_0 = \log 1.9$, we have $e^{-\frac{1}{3} \lambda_0 (l-k)} 2^{53(l-k)/200} \leq e^{-(l-k)/34}$. So the sum becomes

$$\sum_k \sum_l \frac{16M l e^{l/42} e^{-(l-k)/34} (1.9)^{-k/2}}{d'_0 \varepsilon \sqrt{|x - \hat{T}^k 0|}} = \sum_k \frac{16M e^{k/34} (1.9)^{-k/2}}{d'_0 \varepsilon \sqrt{|x - \hat{T}^k 0|}} \sum_l l e^{-l/34} e^{l/42}$$

and both series converge.

ρ_1 has bounded variation. Note that $p_{l,j}^{(i)}$ is constant on $\Delta_{l,j}^{(i)}$. We set

$$\rho_1 = \sum_{\Delta_{l,j}^{(i)}: l_j \geq p_{l,j}^{(i)}} \frac{\varphi \circ \pi_{l,j}^{(i)-1}}{\pi' \circ \pi_{l,j}^{(i)-1}}.$$

Equation (35) and equation (28) of §4.5 imply that

$$\left| \frac{(\pi_{l,j}^{(i)-1})'(x)}{(\pi_{l,j}^{(i)-1})'(y)} - 1 \right| \leq \tilde{C},$$

which in turns yields the bound we need to estimate the variation:

$$\left| \frac{(\pi_{l,j}^{(i)-1})''(x)}{(\pi_{l,j}^{(i)-1})'(x)} \right| \leq \tilde{C}. \quad (37)$$

Let $\bigvee_J f$ denote the variation of a function on the interval J . Recall that the regularity functional on the tower is given by $\|f\|_{\Delta_{l,j}^{(i)}} = \sup_{\Delta_{l,j}^{(i)}} |f'/f| e^{-\xi l}$ for any $f \in X$. Thus

$$\bigvee_{\Delta_{l,j}^{(i)}} f = \int_{\Delta_{l,j}^{(i)}} |f'| dm \leq \|f\|_{\Delta_{l,j}^{(i)}} e^{\xi l} \int_{\Delta_{l,j}^{(i)}} |f| dm. \quad (38)$$

We now estimate the variation of ρ_1 on I :

$$\bigvee_I \rho_1 \leq \sum_{\Delta_{l,j}^{(i)}: l_j \geq p_{l,j}^{(i)}} \bigvee_{\pi(\Delta_{l,j}^{(i)})} \varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})' + 2 \left| \frac{\varphi \circ \pi_{l,j}^{(i)-1}}{\pi' \circ \pi_{l,j}^{(i)-1}} \right|_{\infty}.$$

Since the $\Delta_{l,j}^{(i)}$ in the sum are free, we estimate the second term using

$$\left| \frac{\varphi \circ \pi_{l,j}^{(i)-1}}{\pi' \circ \pi_{l,j}^{(i)-1}} \right|_{\infty} \leq \frac{M e^{-\frac{1}{3}\lambda_0 l}}{\lambda^l c'_0 c_0 \delta}. \quad (39)$$

We estimate the first term one $\Delta_{l,j}^{(i)}$ at a time:

$$\begin{aligned} \bigvee_{\pi(\Delta_{l,j}^{(i)})} \varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})' &= \int_{\pi(\Delta_{l,j}^{(i)})} |(\varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})')'| dm \\ &\leq \int_{\pi(\Delta_{l,j}^{(i)})} |\varphi' \circ \pi_{l,j}^{(i)-1} \cdot ((\pi_{l,j}^{(i)-1})')^2| dm \\ &\quad + \int_{\pi(\Delta_{l,j}^{(i)})} |\varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})''| dm \\ &\leq \frac{e^{-\frac{1}{3}\lambda_0 l}}{c'_0 c_0 \delta} \int_{\Delta_{l,j}^{(i)}} |\varphi'| dm + \tilde{C} \int_{\Delta_{l,j}^{(i)}} \varphi dm, \end{aligned}$$

where we have used equation (37) for the second term in the last line. We use equation (38) for the first term in the last line to obtain

$$\bigvee_{\pi(\Delta_{l,j}^{(i)})} \varphi \circ \pi_{l,j}^{(i)-1} \cdot (\pi_{l,j}^{(i)-1})' \leq \frac{M e^{(-\frac{1}{3}\lambda_0 + \xi)l}}{c'_0 c_0 \delta} \int_{\Delta_{l,j}^{(i)}} \varphi dm + \tilde{C} \int_{\Delta_{l,j}^{(i)}} \varphi dm. \quad (40)$$

Now putting together equations (39) and (40) we conclude that

$$\bigvee_I \rho_1 \leq \sum_{\Delta_{l,j}^{(i)}: l_j \geq p_{l,j}^{(i)}} \frac{M e^{(-\frac{1}{3}\lambda_0 + \xi)l}}{c'_0 c_0 \delta} \int_{\Delta_{l,j}^{(i)}} \varphi dm + \tilde{C} \int_{\Delta_{l,j}^{(i)}} \varphi dm + 2 \frac{M e^{-\frac{1}{3}\lambda_0 l}}{c'_0 c_0 \delta \lambda^l}.$$

This sum is finite since on each level of the tower, there are only finitely many $\Delta_{l,j}^{(i)}$ which are free and it is only these $\Delta_{l,j}^{(i)}$ which we are summing over. The number of such pieces on a level l of the tower has been shown in equation (34) to be bounded by $8l2^{53l/200}$. Using this estimate, the fact that $\lambda \geq e^{-\frac{1}{42}}$ and the observation that ξ is much smaller than λ_0 , the above series is finite and so ρ_1 has bounded variation.

These estimates show that the density ψ has the form claimed in the statement of Theorem 2.3. Moreover, equation (36) allows us to write $\psi_s \leq g$, where $g \in L^1(\hat{I})$ and g is independent of s in the sequence of holes.

Since the upper and lower bounds on ψ_s are uniform in s , we may conclude that the sequence $\{\mu_s\}$ has a subsequence $\{\mu_{s_k}\}$ which converges weakly to a measure ν that is

bounded away from zero on $[1 - a, 1]$. The limit measure ν cannot be singular because of the uniform upper bound we derived for the sequence $\{\psi_s\}$. Also, since by equation (32) $\lambda_s \rightarrow 1$ as $s \rightarrow 0$, we have for any Borel subset A of $[-1, 1]$,

$$\begin{aligned} \nu(\hat{T}^{-1}A) &= \lim_{k \rightarrow \infty} \mu_{s_k}(\hat{T}^{-1}A) = \lim_{k \rightarrow \infty} \mu_{s_k}(T_{s_k}^{-1}(A \setminus H_{s_k})) \\ &= \lim_{k \rightarrow \infty} \lambda_{s_k} \mu_{s_k}(A \setminus H_{s_k}) = \nu(A) \end{aligned}$$

so that ν is an absolutely continuous invariant measure for \hat{T} with density bounded away from zero on $[1 - a, 1]$. But there is only one such measure for \hat{T} ([Y1]). This implies that the entire sequence $\{\mu_s\}$ converges to the unique ergodic a.c.i.m. for \hat{T} as $m(H_s) \rightarrow 0$.

Acknowledgement. The author is grateful to L.-S. Young for many helpful discussions and guidance during the writing of this paper.

REFERENCES

- [BC] M. Benedicks and L. Carleson. On iterates of $1 - ax^2$ on $(-1, 1)$. *Ann. Math.* **122** (1985), 1–25.
- [BCh] H. van den Bedem and N. Chernov. Expanding maps of an interval with holes. *Ergod. Th. & Dynam. Sys.* **22** (2002), 637–654.
- [BK] V. Baladi and G. Keller. Zeta functions and transfer operators for piecewise monotonic transformations. *Comm. Math. Phys.* **127** (1990), 459–477.
- [BY] M. Benedicks and L.-S. Young. Markov extensions and decay of correlations for certain Hénon maps. *Astérisque* **261** (2000), xi, 13–56.
- [C1] N. N. Čencova. A natural invariant measure on Smale’s horseshoe. *Soviet Math. Dokl.* **23** (1981), 87–91.
- [C2] N. N. Čencova. Statistical properties of smooth Smale horseshoes. *Mathematical Problems of Statistical Mechanics and Dynamics*. Ed. R. L. Dobrushin. Reidel, Dordrecht, 1986, pp. 199–256.
- [Ch1] N. Chernov. Statistical properties of piecewise smooth hyperbolic systems in high dimensions. *Discr. Cont. Dynam. Syst.* **5** (1999), 425–448.
- [Ch2] N. Chernov. Decay of correlations and dispersing billiards. *J. Stat. Phys.* **94** (1999), 513–556.
- [Ch3] N. Chernov. Sinai billiards under small external forces. *Ann. Henri Poincaré* **2** (2001), 197–236.
- [CM1] N. Chernov and R. Markarian. Ergodic properties of Anosov maps with rectangular holes. *Bol. Soc. Bras. Mat.* **28** (1997), 271–314.
- [CM2] N. Chernov and R. Markarian. Anosov maps with rectangular holes. Nonergodic cases. *Bol. Soc. Bras. Mat.* **28** (1997), 315–342.
- [CMS1] P. Collet, S. Martínez and B. Schmitt. The Yorke–Pianigiani measure and the asymptotic law on the limit Cantor set of expanding systems. *Nonlinearity* **7** (1994), 1437–1443.
- [CMS2] P. Collet, S. Martínez and B. Schmitt. Quasi-stationary distribution and Gibbs measure of expanding systems. *Instabilities and Nonequilibrium Structures V*. Eds. E. Tirapegui and W. Zeller. Kluwer, Dordrecht, 1996, pp. 205–219.
- [CMT1] N. Chernov, R. Markarian and S. Troubetsky. Conditionally invariant measures for Anosov maps with small holes. *Ergod. Th. & Dynam. Sys.* **18** (1998), 1049–1073.
- [CMT2] N. Chernov, R. Markarian and S. Troubetsky. Invariant measures for Anosov maps with small holes. *Ergod. Th. & Dynam. Sys.* **20** (2000), 1007–1044.
- [D] M. Demers. Markov extensions for dynamical systems with holes: an application to expanding maps of the interval. *Israel J. Math.* **146** (2005), 189–221.
- [KL] G. Keller and C. Liverani. Stability of the spectrum for transfer operators. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **28** (1999), 141–152.
- [LiM] C. Liverani and V. Maume-Deschamps. Lasota–Yorke maps with holes: conditionally invariant probability measures and invariant probability measures on the survivor set. *Ann. Henri Poincaré Prob. Stat.* **39** (2003), 385–412.

- [LM] A. Lopes and R. Markarian. Open billiards: Cantor sets, invariant and conditionally invariant probabilities. *SIAM J. Appl. Math.* **56** (1996), 651–680.
- [PY] G. Pianigiani and J. Yorke. Expanding maps on sets which are almost invariant: decay and chaos. *Trans. Amer. Math. Soc.* **252** (1979), 351–366.
- [R] P. A. Richardson, Jr. Natural measures on the unstable and invariant manifolds of open billiard dynamical systems. *Doctoral Dissertation*, Department of Mathematics, University of North Texas, 1999.
- [WY1] Q. Wang and L.-S. Young. Strange attractors with one direction of instability. *Comm. Math. Phys.* **218** (2001), 1–97.
- [WY2] Q. Wang and L.-S. Young. Nonuniformly expanding one-dimensional maps. *Preprint*, 2005.
- [Y1] L.-S. Young. Decay of correlations for certain quadratic maps. *Comm. Math. Phys.* **146** (1992), 123–138.
- [Y2] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. Math.* **147** (1998), 585–650.
- [Y3] L.-S. Young. Recurrence times and rates of mixing. *Israel J. Math.* **110** (1999), 153–188.